

Astérisque

PASCAL AUSCHER

PHILIPPE TCHAMITCHIAN

**Square root problem for divergence operators
and related topics**

Astérisque, tome 249 (1998)

<http://www.numdam.org/item?id=AST_1998__249__R1_0>

© Société mathématique de France, 1998, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ASTÉRIQUE 249

**SQUARE ROOT PROBLEM
FOR DIVERGENCE OPERATORS
AND RELATED TOPICS**

**Pascal Auscher
Philippe Tchamitchian**

Société Mathématique de France 1998

Pascal Auscher

LAMFA, Université de Picardie Jules-Verne, 33, rue Saint-Leu,
F-80039 Amiens Cedex 1.

E-mail : `auscher@mathinfo.u-picardie.fr`

Philippe Tchamitchian

Faculté des Sciences et Techniques de Saint-Jérôme, Université d'Aix-Marseille III,
Avenue Escadrille Normandie-Niemen, F-13397 Marseille Cedex 20,
and LATP, CNRS, UMR 6632.

E-mail : `tchamphi@math.u-3mrs.fr`

1991 Mathematics Subject Classification. — Primary: 35J15, 47A60, 42B25. Secondary: 42B20, 35K10, 47F05, 46G20.

Key words and phrases. — Second order complex elliptic operators, maximal accretive operators, H^∞ -functional calculus, multilinear expansions, Gaussian estimates, Littlewood-Paley-Stein quadratic functionals, Carleson measures, absolutely bounded mean oscillation, Riesz transforms, Calderón-Zygmund operators.

SQUARE ROOT PROBLEM FOR DIVERGENCE OPERATORS AND RELATED TOPICS

Pascal Auscher, Philippe Tchamitchian

Abstract. — We present in this work recent progress on the square root problem of Kato for differential operators in divergence form on $\mathbb{R}^n$. We discuss topics on functional calculus, heat and resolvent kernel estimates, square function estimates and Carleson measure estimates for square roots. In the first chapter, we show in a quantitative way how the theorems of Aronson-Nash and of De Giorgi are equivalent. In the central chapters, we take advantage of recent development in functional calculus and in harmonic analysis to propose a new point of view on Kato's problem which allows us to unify previous results and extend them. In the last chapter we study the associated Riesz transforms, their relation to Calderón-Zygmund operators and their behavior on L^p -spaces.

Résumé (Problème de la racine carrée pour les opérateurs sous forme divergence et sujets connexes). — Ce travail a pour thème principal le problème de Kato concernant la racine carrée des opérateurs différentiels elliptiques sous forme divergence dans $\mathbb{R}^n$. Pour mener à bien cette étude, nous nous intéressons à des questions relatives au calcul fonctionnel, aux estimations de noyaux, aux fonctionnelles quadratiques et aux mesures de Carleson associées aux racines carrées. Dans le premier chapitre, nous montrons en un sens précis comment les théorèmes d'Aronson-Nash et de De Giorgi sont équivalents. Dans les deux chapitres centraux, nous tirons parti de développements récents sur le calcul fonctionnel et en analyse harmonique pour proposer un nouveau point de vue sur le problème de Kato qui permet d'unifier les résultats antérieurs et de les généraliser. Enfin, dans le dernier chapitre, nous étudions les transformées de Riesz associées, leur relation aux opérateurs de Calderón-Zygmund et leur comportement sur les espaces L^p .

Contents

Introduction	1
Preliminaries	7
0.1. Maximal accretive operators and their functional calculus	7
0.2. Maximal accretive operators and forms	9
0.3. The square root problem from an abstract point of view	12
0.4. The square root problem for differential operators on $\mathbb{R}^n$	14
0.5. The square root problem and perturbation theory	16
0.6. Connections between second and higher order operators	18
0.7. Perturbations with lower order terms	19
0.8. Change of variables	22
0.9. Further comments	25
1. Gaussian estimates	27
1.1. Introduction	27
1.2. Gaussian estimates for second order operators	28
1.3. A counterexample	32
1.4. The Dirichlet property and the Gaussian property	33
1.4.1. Dirichlet integrals and the Dirichlet property	33
1.4.2. From the Gaussian property to the Dirichlet property	35
1.4.3. From the Dirichlet property to the Gaussian property	37
1.4.4. Equivalence of the local properties	44
1.4.5. Inhomogeneous operators	44
1.4.6. Stability of (D) and proof of Theorem 6	45
1.4.7. Gradient estimates on the heat kernel	48
1.5. Further consequences of the Gaussian property	52
1.5.1. Green kernel estimates	52
1.5.2. L^p estimates	53
1.5.3. The conservation property	55
1.6. Analytic perturbation	56
1.7. Higher order elliptic operators	58
2. Quadratic functionals, Carleson measures and square roots of differential operators	65
2.1. Introduction	65
2.2. Classical quadratic functionals	65
2.2.1. A review	65

2.2.2. Application to the one dimensional square root problem	69
2.2.3. Quadratic functionals of weakly regular families	70
2.2.4. Square roots of operators with special structures	76
2.3. Counterexamples	77
2.4. Quadratic functionals of irregular families	80
2.5. The heart of the matter	82
2.5.1. Characterizations of (K) in terms of quadratic functionals	83
2.5.2. Characterizations of (K) in terms of Carleson measures	84
2.6. Miscellaneous results	86
2.6.1. The inequality $(K)_{loc}$	86
2.6.2. Localization principles	87
3. Positive answers to the square root problem	91
3.1. Introduction	91
3.2. The class S and the inequality (K)	92
3.3. Applications	97
3.3.1. Structure conditions: Tb -type results	97
3.3.2. Functions of absolutely bounded mean oscillations	100
3.3.3. Perturbations in the BMO topology	103
3.3.4. Open questions relative to the class S	105
3.4. The real symmetric case	105
3.5. A variation on regularity conditions	110
4. Square roots of differential operators, singular integrals and L^p theory	113
4.1. Introduction	113
4.2. Standard factorization of $L^{1/2}$ and Riesz transforms associated to L	115
4.2.1. Generalities	115
4.2.2. Negative results on Riesz transforms	119
4.2.3. Positive results on Riesz transforms	121
4.3. Non standard factorization of $L^{1/2}$ by a Calderón-Zygmund operator	125
4.4. Further L^p estimates and invertibility of $L^{1/2}$	130
4.5. Coefficients depending on one variable	132
4.6. Local L^p estimates	134
4.7. Miscellaneous results	136
4.7.1. Perturbation theory	136
4.7.2. Extensions to higher order operators	137
4.7.3. L^p theory and bilipschitz changes of variables	137
4.7.4. Estimates on Morrey spaces and application	138
A. The space $ABMO$	141
A.1. Proof of Proposition 15 of Chapter 3	141
A.2. Littlewood-Paley characterization of $ABMO$	143
A.3. A subclass of $ABMO$	144
A.4. Further examples	146

B. Coefficients depending on one variable	149
C. Improved constants	155
D. Reduction of dimension principle	161
Bibliography	167

INTRODUCTION

The main goal of this memoir is the study of square roots of divergence form differential operators $L = -\operatorname{div}(A\nabla)$ on $\mathbb{R}^n$, $n \geq 2$, where $A(x)$ is a matrix with complex-valued bounded entries and satisfying a uniform ellipticity condition. This operator is defined using the theory of maximal accretive operators on $L^2(\mathbb{R}^n)$ via the associated variational form on the Sobolev space $H^1(\mathbb{R}^n)$. We denote by $L^{1/2}$ the square root of L .

We are interested in the following questions.

Question 1. — *Does the domain of $L^{1/2}$ agree with $H^1(\mathbb{R}^n)$?*

When L is a selfadjoint operator or has smooth coefficients (e.g., Lipschitz), this is true. However, the answer is not known in general.

Question 2. — *If A_0 is such that the domain of $L_0^{1/2}$ is $H^1(\mathbb{R}^n)$, where $L_0 = -\operatorname{div}(A_0\nabla)$, is the map $A \rightarrow L^{1/2}$, valued in $\mathcal{B}(H^1(\mathbb{R}^n), L^2(\mathbb{R}^n))$, complex analytic at A_0 ?*

Classical complex function theory tells us that analyticity of a Taylor expansion follows from boundedness in complex balls. This is the same principle that partly justifies the use of complex-valued coefficients here.

Question 3. — *If L is such that the domain of $L^{1/2}$ agrees with $H^1(\mathbb{R}^n)$, what can we say about L^p -boundedness, $p \neq 2$, of the Riesz transforms associated to L , namely $\nabla L^{-1/2}$? More generally, how do $\|L^{1/2}f\|_p$ and $\|\nabla f\|_p$ compare?*

In this work we bring new answers to the first two questions and, under a technical hypothesis that the kernel of the semigroup generated by $-L$ has Gaussian upper bounds and regularity, we completely elucidate the third one. Some results were announced in [15].