

373

ASTÉRISQUE

2015

EQUILIBRIUM STATES IN NEGATIVE CURVATURE

Frédéric PAULIN, Mark POLLICOTT and Barbara SCHAPIRA

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Astérisque est un périodique de la Société Mathématique de France.

Numéro 373, 2015

Comité de rédaction

Ahmed ABBES	Damien GABORIAU
Viviane BALADI	Michael HARRIS
Gérard BESSON	Fabrice PLANCHON
Laurent BERGER	Pierre SCHAPIRA
Philippe BIANE	Bertrand TOËN
Hélène ESNAULT	
Éric VASSEROT (dir.)	

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 75 € (\$ 112)

Abonnement Europe : 530 €, hors Europe : 569 € (\$ 853)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2015

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0303-1179

ISBN 978-2-85629-818-3

Directeur de la publication : Marc Peigné

373

ASTÉRISQUE

2015

EQUILIBRIUM STATES IN NEGATIVE CURVATURE

Frédéric PAULIN, Mark POLLICOTT and Barbara SCHAPIRA

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Frédéric Paulin

Département de Mathématique

Bât. 425

UMR 8628 CNRS

Université Paris-Sud

91405 Orsay Cedex (France)

`frederic.paulin@math.u-psud.fr`

Mark Pollicott

Mathematics Institute

University of Warwick

Coventry, CV4 7AL, UK

`mpollic@maths.warwick.ac.uk`

Barbara Schapira

LAMFA, UMR 7352 CNRS

UFR des Sciences

Université Picardie Jules Verne

33 rue Saint Leu

80000 Amiens (France)

`barbara.schapira@u-picardie.fr`

Classification mathématique par sujet (2000). — 37D35, 53D25, 37D40, 37A25, 37C35, 53C12.

Mots-clefs. — Geodesic flow, negative curvature, Gibbs state, periods, orbit counting, Patterson density, pressure, variational principle, strong unstable foliation.

EQUILIBRIUM STATES IN NEGATIVE CURVATURE

by Frédéric PAULIN, Mark POLLICOTT and Barbara SCHAPIRA

Abstract. — With their origin in thermodynamics and symbolic dynamics, Gibbs measures are crucial tools to study the ergodic theory of the geodesic flow on negatively curved manifolds. We develop a framework (through Patterson-Sullivan densities) allowing us to get rid of compactness assumptions on the manifold, and prove many existence, uniqueness and finiteness results of Gibbs measures. We give many applications, to the Variational Principle, the counting and equidistribution of orbit points and periods, the unique ergodicity of the strong unstable foliation and the classification of Gibbs densities on some Riemannian covers.

Résumé (États d'équilibre en courbure négative.) — Les mesures de Gibbs, utilisées d'abord en thermodynamique et en dynamique symbolique, sont des outils cruciaux pour l'étude de la théorie ergodique des flots géodésiques des variétés de courbure strictement négative. Nous introduisons (via les densités de Patterson-Sullivan) un cadre qui permet de s'affranchir d'hypothèses de compacité sur la variété, et nous démontrons de nombreux résultats d'existence, d'unicité et de finitude des mesures de Gibbs. Nous en donnons moult applications, au principe variationnel, au comptage et à l'équidistribution des points d'orbites et des périodes, à l'unique ergodicité du feuilletage fortement instable, et à la classification des mesures de Gibbs sur certains revêtements riemanniens.

CONTENTS

1. Introduction	1
2. Background on negatively curved manifolds	13
2.1. Uniform local Hölder-continuity	14
2.2. Boundary at infinity, isometries and the Busemann cocycle	15
2.3. The geometry of the unit tangent bundle	18
2.4. Geodesic flow, (un)stable foliations and the Hamenstädt distances	21
2.5. Some exercises in hyperbolic geometry	25
2.6. Pushing measures by branched covers	28
3. A Patterson-Sullivan theory for Gibbs states	31
3.1. Potential functions and their periods	31
3.2. The Poincaré series and the critical exponent of (Γ, F)	34
3.3. The Gibbs cocycle of (Γ, F)	39
3.4. The potential gap of (Γ, F)	44
3.5. The crossratio of (Γ, F)	48
3.6. The Patterson densities of (Γ, F)	51
3.7. The Gibbs states of (Γ, F)	57
3.8. The Gibbs property of Gibbs states	59
3.9. Conditional measures of Gibbs states on strong (un)stable leaves	66
4. Critical exponent and Gurevich pressure	75
4.1. Counting orbit points and periodic geodesics	75
4.2. Logarithmic growth of the orbital counting functions	77
4.3. Equality between critical exponent and Gurevich pressure	85
4.4. Critical exponent of Schottky semigroups	87
5. A Hopf-Tsuji-Sullivan-Roblin theorem for Gibbs states	105
5.1. Some geometric notation	105
5.2. The Hopf-Tsuji-Sullivan-Roblin theorem for Gibbs states	108
5.3. Uniqueness of Patterson densities and of Gibbs states	118
6. Thermodynamic formalism and equilibrium states	123
6.1. Measurable partitions and entropy	125
6.2. Proof of the variational principle	127

7. The Liouville measure as a Gibbs measure	143
7.1. The Hölder-continuity of the (un)stable Jacobian	146
7.2. Absolute continuity of the strong unstable foliation	150
7.3. The Liouville measure as an equilibrium state	152
7.4. The Liouville measure satisfies the Gibbs property	155
7.5. Conservative Liouville measures are Gibbs measures	159
8. Finiteness and mixing of Gibbs states	165
8.1. Babillot's mixing criterion for Gibbs states	165
8.2. A finiteness criterion for Gibbs states	168
9. Growth and equidistribution of orbits and periods	175
9.1. Convergence of measures on the square product	175
9.2. Counting orbit points of discrete groups	183
9.3. Equidistribution and counting of periodic orbits of the geodesic flow	185
9.4. The case of infinite Gibbs measure	196
10. The ergodic theory of the strong unstable foliation	199
10.1. Quasi-invariant transverse measures	199
10.2. Quasi-invariant measures on the space of horospheres	203
10.3. Classification of quasi-invariant transverse measures for $\mathcal{W}^{\text{su}}$	206
11. Gibbs states on Galois covers	233
11.1. Improving Mohsen's shadow lemma	234
11.2. The Fatou-Roblin radial convergence theorem	238
11.3. Characters and critical exponents	240
11.4. Characters and Patterson densities	246
11.5. Characters and the Hopf-Tsuji-Sullivan-Roblin theorem	249
11.6. Galois covers and critical exponents	253
11.7. Classification of ergodic Patterson densities on nilpotent covers	257
List of symbols	261
Index	265
Bibliography	271

CHAPTER 1

INTRODUCTION

With their thermodynamic origin, Gibbs measures (or states) are very useful in symbolic dynamics over finite alphabets (see for instance [140], [158], [83]). Sinai, Bowen and Ruelle introduced them in hyperbolic dynamics (which, via coding theory, has strong links to symbolic dynamics), in particular in order to study the weighted distribution of periodic orbits and the equilibrium states for given potentials (see for instance [149], [25], [27], [119]). For instance, this allows the dynamical analysis of the geodesic flow of negatively curved Riemannian manifolds, provided its non-wandering set is compact. As a first step to venture beyond the compact case, Gibbs measures in symbolic dynamics over countable alphabets have been developed by Sarig [142], [144]. But since no coding theory which does not lose geometric information is known for general non-compact manifolds, this methodology is not well adapted to the non-compact case. Note that a coding-free approach of transfer operators, in particular via improved spectral methods, was put in place by Liverani, Baladi, Gouëzel and others (see for instance, restricting the references to the case of flows, [93], [34], [153], [154], [12], [60], as well as [5] for an example in a non locally homogeneous and non-compact situation). In this text, we construct and study geometrically Gibbs measures for the geodesic flow of negatively curved Riemannian manifolds, without compactness assumptions.

By considering the action on a universal Riemannian cover of its covering group, this study is strongly related to the (weighted) distribution of orbits of discrete groups of isometries of negatively curved simply connected Riemannian manifolds. After work of Huber and Selberg (in particular through his trace formula) in constant curvature, Margulis (see for instance [99]) made a breakthrough, albeit in the unweighted lattice case, to give the precise asymptotic growth of the orbits. Patterson [121] and Sullivan [151], in the surface and constant curvature case respectively, have introduced a nice approach using measures at infinity, giving in particular a nice construction of the measure of maximal entropy in the cocompact case. The Patterson-Sullivan theory extends to variable curvature and non lattice case, and an optimal reference (in the unweighted case) is due to Roblin [134].

After preliminary work of Hamenstädt [66] on Gibbs cocycles, of Ledrappier [91], Coudène [38] and Schapira [147] using a slightly different approach, and of Mohsen [103] (their contributions will be explained as the story unfolds), the aim of this book is to develop an optimal theory of Gibbs measures for the dynamical study of geodesic flows in general negatively curved Riemannian manifolds, and of weighted distribution of orbits of general discrete groups of isometries of negatively curved simply connected Riemannian manifolds.

Let us give a glimpse of our results, starting by describing the players. Let M be a complete connected Riemannian manifold with pinched negative curvature. For simplicity in this introduction, we assume that the derivatives of the sectional curvature of M are uniformly bounded. Let $F : T^1M \rightarrow \mathbb{R}$ be a Hölder-continuous map, called a *potential*, which is going to help us define the various weights. In order to simplify the exposition of this introduction, we assume here that F is invariant by the antipodal map $v \mapsto -v$ (see the main body of the text for complete statements). Let $p : \widetilde{M} \rightarrow M$ be a universal Riemannian covering map, with covering group Γ and sphere at infinity $\partial_\infty \widetilde{M}$, and let $\widetilde{F} = F \circ p$. We make no compactness assumption on M : we only assume Γ to be non-elementary (that is, non virtually nilpotent). (In the body of the text, we will also allow M to be a good orbifold, hence Γ to have torsion.) Let $\phi = (\phi_t)_{t \in \mathbb{R}}$ be the geodesic flow on T^1M and $\widetilde{\phi} = (\widetilde{\phi}_t)_{t \in \mathbb{R}}$ the one on $T^1\widetilde{M}$. For all $x, y \in \widetilde{M}$, let us define

$$\int_x^y \widetilde{F} = \int_0^{d(x,y)} \widetilde{F}(\widetilde{\phi}_t v) dt$$

where v is the unit tangent vector at x to a geodesic from x through y . For every periodic orbit g of ϕ , let $\mathcal{L}_g$ be the Lebesgue measure along g and $\int_g F = \mathcal{L}_g(F)$ the *period* of g for the potential F .

We will study three numerical invariants of the weighted dynamics.

- Let $x, y \in \widetilde{M}$, and $c > 0$ large enough. The *critical exponent* of (Γ, F) is an exponential growth rate of the orbit points of Γ weighted by the potential F :

$$\delta_{\Gamma, F} = \lim_{t \rightarrow +\infty} \frac{1}{t} \log \sum_{\substack{\gamma \in \Gamma \\ t-c \leq d(x, \gamma y) \leq t}} e^{\int_x^{\gamma y} \widetilde{F}}.$$

- For every $t \geq 0$, let $\mathcal{P}\text{er}(t)$ be the set of periodic orbits of ϕ with length at most t , and $\mathcal{P}\text{er}'(t)$ its subset of primitive ones. Let W be a relatively compact open subset of T^1M meeting the non-wandering set of ϕ , and $c > 0$ large enough. The *Gurevich pressure* of (M, F) is an exponential growth rate of the closed geodesics of M weighted by the potential F :

$$P_{\text{Gur}}(M, F) = \lim_{t \rightarrow +\infty} \frac{1}{t} \log \sum_{\substack{g \in \mathcal{P}\text{er}(t) - \mathcal{P}\text{er}(t-c) \\ g \cap W \neq \emptyset}} e^{\int_g \widetilde{F}}.$$

The restriction for the periodic orbits under consideration to meet a given compact set is necessary: For instance if M is an infinite cyclic cover of a compact manifold,