

quatrième série - tome 52 fascicule 4 juillet-août 2019

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Siddhartha SAHI & Hadi SALMASIAN

Quadratic Capelli operators and Okounkov polynomials

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Patrick BERNARD

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE

de 1883 à 1888 par H. DEBRAY

de 1889 à 1900 par C. HERMITE

de 1901 à 1917 par G. DARBOUX

de 1918 à 1941 par É. PICARD

de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} mars 2019

P. BERNARD

D. HARARI

S. BOUCKSOM

A. NEVES

R. CERF

J. SZEFTEL

G. CHENEVIER

S. VŨ NGỌC

Y. DE CORNULIER

A. WIENHARD

A. DUCROS

G. WILLIAMSON

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,

45, rue d'Ulm, 75230 Paris Cedex 05, France.

Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.

annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France

Case 916 - Luminy

13288 Marseille Cedex 09

Tél. : (33) 04 91 26 74 64

Fax : (33) 04 91 41 17 51

email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 420 euros.

Abonnement avec supplément papier :

Europe : 551 €. Hors Europe : 620 € (\$ 930). Vente au numéro : 77 €.

© 2019 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Stéphane Seuret

Périodicité : 6 n^{os} / an

QUADRATIC CAPELLI OPERATORS AND OKOUNKOV POLYNOMIALS

BY SIDDHARTHA SAHI AND HADI SALMASIAN

ABSTRACT. — Let Z be the symmetric cone of $r \times r$ positive definite Hermitian matrices over a real division algebra \mathbb{F} . Then Z admits a natural family of invariant differential operators—the *Capelli operators* C_λ —indexed by partitions λ of length at most r , whose eigenvalues are specializations of Knop-Sahi interpolation polynomials.

In this paper we consider a double fibration $Y \leftarrow X \rightarrow Z$ where Y is the Grassmanian of r -dimensional subspaces of \mathbb{F}^n with $n \geq 2r$. Using this we construct a family of invariant differential operators $D_{\lambda,s}$ on Y that we refer to as *quadratic Capelli operators*. Our main result shows that the eigenvalues of the $D_{\lambda,s}$ are specializations of Okounkov interpolation polynomials.

RÉSUMÉ. — Soit Z le cône symétrique de matrices de tailles $r \times r$ hermitiennes positives sur une algèbre de division réelle \mathbb{F} . Alors Z admet une famille naturelle d'opérateurs différentiels invariants — les *Opérateurs de Capelli* C_λ — indexés par des partitions λ de longueur au plus r , dont les valeurs propres sont des spécialisations de polynômes d'interpolation Knop-Sahi.

Dans cet article, nous considérons une double fibration $Y \leftarrow X \rightarrow Z$ où Y est la variété grassmannienne des sous-espaces de dimension r de \mathbb{F}^n avec $n \geq 2r$. En utilisant cela, nous construisons une famille d'opérateurs différentiels invariants $D_{\lambda,s}$ sur Y que nous appelons opérateurs de Capelli *quadratiques*. Notre résultat principal montre que les valeurs propres des $D_{\lambda,s}$ sont des spécialisations de polynômes d'interpolation Okounkov.

1. Introduction

Let $\mathbb{F} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ be a real division algebra. Fix integers r and n such that $1 \leq r \leq \frac{n}{2}$. Let Y be the Grassmannian of r -dimensional subspaces of \mathbb{F}^n , and let Z be the symmetric cone of $r \times r$ positive definite Hermitian \mathbb{F} -matrices. Then one has a double fibration

$$\begin{array}{ccc} & X & \\ \varphi \swarrow & & \searrow \psi \\ Y & & Z, \end{array}$$

where X is the space of $n \times r$ matrices of \mathbb{F} -rank r . For $x \in X$, $\varphi(x)$ is the column space (or range) of x , while $\psi(x) := x^\dagger x$, where x^\dagger denotes the \mathbb{F} -Hermitian adjoint of x .

One can give another description of the above structure in terms of the groups

$$G_m := \text{GL}_m(\mathbb{F}), \quad K_m := \text{U}_m(\mathbb{F}) := \left\{ g \in G_m : g^\dagger g = I_{m \times m} \right\}.$$

The groups K_n and G_r act on X by matrix multiplication on the left and right respectively, and the maps $X \xrightarrow{\psi} Z$ and $X \xrightarrow{\varphi} Y$ are simply the corresponding quotient maps. Moreover, $X \xrightarrow{\varphi} Y$ is a principal G_r -bundle, while $X \xrightarrow{\psi} Z$ is a fibration whose fibers are isomorphic to the Stiefel manifold K_n/K_{n-r} . Also, since the actions of K_n and G_r on X commute, it follows that G_r acts on Z , and K_n acts on Y . In fact, Y and Z are symmetric spaces for the latter actions. More precisely, we have

$$Y \simeq K_n/(K_r \times K_{n-r}) \text{ and } Z \simeq G_r/K_r.$$

The cone Z is a symmetric space of type A , and admits an important basis of G_r -invariant differential operators C_λ , indexed by partitions $\lambda \in \mathcal{P}_r$, where

$$\mathcal{P}_r := \{(\lambda_1, \dots, \lambda_r) \in \mathbb{Z}^r : \lambda_1 \geq \dots \geq \lambda_r \geq 0\}.$$

The operators C_λ were first studied by the first author in [20], and were referred to as *Capelli operators*. It is known that the spectrum of C_λ is given by specialization of Knop-Sahi type A interpolation polynomials [20], [12], [30].

On the other hand, Y is a compact symmetric space of type BC . In this paper, we use the above double fibration to construct a family of K_n -invariant differential operators $D_{\lambda,s}$ on Y that correspond to the Capelli operators C_λ . We call the operators $D_{\lambda,s}$ the *quadratic Capelli operators* because they are obtained from C_λ by pullback of the quadratic map ψ . Our main result proves that the spectrum of $D_{\lambda,s}$ is given by specialization of the *Okounkov type BC* interpolation polynomials $P_\lambda(x; \tau, \alpha)$ (see [14, Sec. 5.3] and [19]).

To describe our main result precisely, we begin by introducing some notation. Set $K := K_n$ and $M := K_r \times K_{n-r} \subset K$, so that $Y \simeq K/M$. The group K acts by left translation on $C^\infty(Y)$, the space of complex-valued smooth functions on Y . The operators $D_{\lambda,s}$ leave the subspace $C^\infty(Y)_{K\text{-finite}}$ of K -finite vectors invariant. By standard results from the theory of compact symmetric spaces (for example, see [5, Chap. V]), $C^\infty(Y)_{K\text{-finite}}$ decomposes as a multiplicity-free direct sum of irreducible M -spherical K -modules, which are naturally parametrized by partitions $\mu \in \mathcal{P}_r$. Our next goal is to describe this parametrization. Let \mathfrak{k} and \mathfrak{m} denote the Lie algebras of K and M . Fix a Cartan decomposition $\mathfrak{k} = \mathfrak{m} \oplus \mathfrak{p}$. Let $\mathfrak{a} \subseteq \mathfrak{p}$ be a Cartan subspace, and let \mathfrak{h} be a Cartan subalgebra of \mathfrak{k} such that $\mathfrak{a} \subseteq \mathfrak{h}$. Then $\mathfrak{h} = \mathfrak{t} \oplus \mathfrak{a}$, where $\mathfrak{t} := \mathfrak{h} \cap \mathfrak{m}$. Set $\mathfrak{k}_\mathbb{C} := \mathfrak{k} \otimes_\mathbb{R} \mathbb{C}$, $\mathfrak{h}_\mathbb{C} := \mathfrak{h} \otimes_\mathbb{R} \mathbb{C}$, $\mathfrak{a}_\mathbb{C} := \mathfrak{a} \otimes_\mathbb{R} \mathbb{C}$, and $\mathfrak{t}_\mathbb{C} := \mathfrak{t} \otimes_\mathbb{R} \mathbb{C}$. The restricted root system $\Sigma := \Sigma(\mathfrak{k}_\mathbb{C}, \mathfrak{a}_\mathbb{C})$ is of type BC_r . We choose a positive system $\Sigma^+ \subset \Sigma$ and a basis e_1, \dots, e_r of $\mathfrak{a}_\mathbb{C}^*$ such that the multiplicity m_α of every $\alpha \in \Sigma^+$ is given in terms of n, r , and $d := \dim \mathbb{F}$ in Table 1 below.

α	$e_i, 1 \leq i \leq r$	$e_i \pm e_j, 1 \leq i < j \leq r$	$2e_i, 1 \leq i \leq r$
m_α	$d(n - 2r)$	d	$d - 1.$

Table 1.

We also choose a positive system for the root system $\Delta := \Delta(\mathfrak{k}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}})$ which is compatible with Σ^+ . Let $\tilde{\mu} \in \mathfrak{h}_{\mathbb{C}}^*$. By the Cartan-Helgason theorem, $\tilde{\mu}$ is the highest weight of an irreducible M -spherical K -module if and only if

$$(1) \quad \tilde{\mu}|_{\mathfrak{t}_{\mathbb{C}}} = 0 \text{ and } \tilde{\mu}|_{\mathfrak{a}_{\mathbb{C}}} = \sum_{i=1}^r 2\mu_i e_i, \text{ where } \mu := (\mu_1, \dots, \mu_r) \in \mathcal{P}_r.$$

REMARK 1.1. – Assume that $\mathbb{F} = \mathbb{R}$. Then K is disconnected, and if K° denotes the connected component of identity of K , then $M \cap K^\circ$ is also disconnected. Therefore the Cartan-Helgason theorem as stated for instance in [5, Cor. V.4.2] does not apply immediately to the case $\mathbb{F} = \mathbb{R}$. However, one can use the refinement of the Cartan-Helgason theorem for the pair $(K^\circ, M \cap K^\circ)$, given in [4, Sec. 12.3.2], as well as the description of irreducible representations of K in terms of irreducible representations of K° , given in [4, Thm 5.5.23], to obtain the condition (1).

From now on, we denote the M -spherical K -module with highest weight $\tilde{\mu}$ satisfying (1) by V_μ . Therefore as K -modules,

$$C^\infty(Y)_{K\text{-finite}} \simeq \bigoplus_{\mu \in \mathcal{P}_r} V_\mu.$$

The operator $D_{\lambda,s}$ acts on V_μ by the scalar

$$c_{\lambda,s}(\mu) := \text{HC}(D_{\lambda,s})(\tilde{\mu}|_{\mathfrak{a}_{\mathbb{C}}} + \rho),$$

where $\rho := \frac{1}{2} \sum_{\alpha \in \Sigma^+} \alpha$, $\tilde{\mu}$ is the highest weight of V_μ , and $\text{HC} : \mathbf{D}_K(Y) \rightarrow \mathcal{P}(\mathfrak{a}_{\mathbb{C}}^*)^W$ is the Harish-Chandra homomorphism from the algebra $\mathbf{D}_K(Y)$ of K -invariant differential operators on Y onto the algebra of polynomials on $\mathfrak{a}_{\mathbb{C}}^*$ that are invariant under the action of the restricted Weyl group W .

We now recall the definition of the Okounkov polynomials $P_\lambda(x; \tau, \alpha)$. Let $\mathbb{k} := \mathbb{C}(\tau, \alpha)$ denote the field of rational functions in τ and α . Let $\delta, \mathbb{1} \in \mathcal{P}_r$ and $\varrho_{\tau,\alpha} \in \mathbb{k}^r$ be defined by

$$(2) \quad \delta := (r - 1, \dots, 0), \quad \mathbb{1} := (1, \dots, 1), \quad \varrho_{\tau,\alpha} := \tau\delta + \alpha\mathbb{1}.$$

For $\lambda \in \mathcal{P}_r$, we define $|\lambda| := \sum_{i=1}^r \lambda_i$. Up to a scalar, $P_\lambda(x; \tau, \alpha) \in \mathbb{k}[x_1, \dots, x_r]$ is the unique polynomial of degree $2|\lambda|$, which is invariant under permutations and sign changes of x_1, \dots, x_r , and satisfies

$$P_\lambda(\mu + \varrho_{\tau,\alpha}; \tau, \alpha) = 0$$

for every $\mu \in \mathcal{P}_r$ such that $|\mu| \leq |\lambda|$ and $\mu \neq \lambda$ (for more details, see Section 4).

Recall that $d := \dim \mathbb{F}$. Let $\mathfrak{i} : \mathbb{C}^r \rightarrow \mathfrak{a}_{\mathbb{C}}^*$ be the linear map defined by $\mathfrak{i}(e^i) := 2e_i$ for $1 \leq i \leq r$, where e^1, \dots, e^r are the standard basis vectors of \mathbb{C}^r (therefore $\mathfrak{i}(\mathcal{P}_r)$ is the set of restrictions to $\mathfrak{a}_{\mathbb{C}}$ of highest weights of M -spherical K -modules). Set

$$\varrho := \mathfrak{i}^{-1}(\rho).$$

A simple calculation yields

$$(3) \quad \varrho = (\varrho_1, \dots, \varrho_r) \text{ where } \varrho_i := \frac{dn}{4} - \frac{1}{2} - \frac{d(i-1)}{2} \text{ for every } 1 \leq i \leq r.$$

We are now ready to state our main theorem.