

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

**Tome 150
Fascicule 3**

2022

Gaëtan Chenevier — Statistics for Kneser p -neighbors	473-516
David Leturcq — Bott–Cattaneo–Rossi invariants for long knots in asymptotic homology $\mathbb{R}^3$	517-542
Marian Aprodu, Indranil Biswas, Sorin Dumitrescu & Sebastian Heller — On the monodromy map for logarithmic differential systems	543-568
Sergey Lysenko — Towards canonical representations of finite Heisenberg groups	569-577
Laurent Michel — Spectral asymptotics for the Metropolis algorithm on singular domains	579-623

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Pages 473-623

Sommaire

Gaëtan Chenevier — Statistiques et p -voisins de Kneser	473-516
David Leturcq — Invariants de Bott-Catteno-Rossi des nœuds longs dans les $\mathbb{R}^3$ d'homologie asymptotiques	517-542
Marian Aprodu, Indranil Biswas, Sorin Dumitrescu & Sebastian Heller — Sur la monodromie des systèmes différentiels logarithmiques	543-568
Sergey Lysenko — Vers les représentations canoniques des groupes de Heisenberg finis	569-577
Laurent Michel — Asymptotiques spectrales pour l'algorithme de Metropolis sur des domaines singuliers	579-623

Contents

Gaëtan Chenevier — Statistics for Kneser p -neighbors 473-516

David Leturcq — Bott–Cattaneo–Rossi invariants for long knots in asymptotic homology $\mathbb{R}^3$ 517-542

Marian Aprodu, Indranil Biswas, Sorin Dumitrescu & Sebastian Heller — On the monodromy map for logarithmic differential systems 543-568

Sergey Lysenko — Towards canonical representations of finite Heisenberg groups 569-577

Laurent Michel — Spectral asymptotics for the Metropolis algorithm on singular domains 579-623

STATISTICS FOR KNESER p -NEIGHBORS

WITH AN APPENDIX BY OLIVIER TAÏBI

BY GAËTAN CHENEVIER

ABSTRACT. — Let L and L' be two integral Euclidean lattices in the same genus. We give an asymptotic formula for the number of Kneser p -neighbors of L that are isometric to L' , when the prime p goes to infinity. In the case when L is unimodular, and if we fix furthermore a subgroup $A \subset L$, we also give an asymptotic formula for the number of p -neighbors of L containing A and which are isomorphic to L' . These statements explain numerical observations in [1], [4], and [12].

In the Appendix, O. Taïbi shows, from Arthur's results, how to deduce the existence of global parameters associated to automorphic representations of definite orthogonal groups over the rationals.

Texte reçu le 5 mai 2021, accepté le 19 octobre 2021.

GAËTAN CHENEVIER, CNRS, DMA, École Normale Supérieure, 45 rue d'Ulm, 75230 Paris Cedex 5, France • *E-mail* : gaetan.chenevier@cnrs.fr

Mathematical subject classification (2010). — 11F; 11F46, 11F55, 11F70, 11F72, 11F80, 11G40, 11H06, 11R39, 11Y55, 22C05.

Key words and phrases. — Integral quadratic forms, unimodular lattices, Kneser neighbors, automorphic forms, equidistribution.

During this work, the authors were supported by C.N.R.S. and by the project ANR-19-CE40-0015-02 COLOSS.