

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

CORRECTIONS TO UPPER BOUNDS FOR COURANT-SHARP NEUMANN AND ROBIN EIGENVALUES

Katie Gittins & Corentin Léna

Tome 154
Fascicule 1

2026

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

pages 265-267

Le *Bulletin de la Société Mathématique de France* est un périodique trimestriel
de la Société Mathématique de France.

Fascicule 1, tome 154, mars 2026

Comité de rédaction

Boris ADAMCZEWSKI
Valeria BANICA
Julie DÉSERTI
Gabriel DOSPINESCU
Dorothee FREY

Youness LAMZOURI
Wendy LOWEN
Ludovic RIFFORD
Erwan ROUSSEAU
Béatrice de TILIÈRE

François DAHMANI (Dir.)

Diffusion

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 9
France
commandes@smf.emath.fr

AMS
P.O. Box 6248
Providence RI 02940
USA
www.ams.org

Tarifs

Vente au numéro : 50 € (\$ 75)

Abonnement électronique : 175 € (\$ 262),

avec supplément papier : Europe 266 €, hors Europe 307 € (\$ 460)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Bulletin de la SMF

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 1 44 27 67 99 • Fax : (33) 1 40 46 90 96

bulletin@smf.emath.fr • smf.emath.fr

© Société Mathématique de France 2026

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directrice de la publication : Isabelle GALLAGHER

CORRECTIONS TO
UPPER BOUNDS FOR COURANT-SHARP
NEUMANN AND ROBIN EIGENVALUES

BY KATIE GITTINS & CORENTIN LÉNA

ABSTRACT. — In [2], the number of nodal domains for the eigenfunctions of the Neumann or Robin Laplacian is bounded from above using the classical (Euclidean) Faber–Krahn inequality. However, in an arbitrary Riemannian manifold, this inequality might not hold. We supply the missing arguments in two dimensions and outline a modification of the method, which preserves most of the results, in n dimensions.

RÉSUMÉ (*Corrections à « Majoration des valeurs propres Courant strictes de Neumann et Robin »*). — Dans [2], on obtient une majoration du nombre de domaines nodaux pour une fonction propre du laplacien de Neumann ou de Robin à l'aide de l'inégalité de Faber-Krahn classique (euclidienne). Cette inégalité n'est toutefois pas nécessairement vérifiée dans une variété riemannienne quelconque. La présente note donne les arguments manquants en deux dimensions et indique à grands traits une modification de la méthode en dimension n qui préserve l'essentiel des résultats.

The mapping F introduced in Equation (6) on page 105 should instead be defined from the cylinder $\mathcal{C}_L := (\mathbb{R}/(L\mathbb{Z})) \times \mathbb{R}$ to \mathbb{R}^2 . The set V^R , defined at the bottom of page 109 through a preimage by F , is, therefore, an open set in \mathcal{C}_L

Texte reçu le 10 avril 2025, accepté le 26 août 2025.

KATIE GITTINS, Department of Mathematical Sciences, Durham University, Mathematical Sciences and Computer Science Building, Upper Mountjoy Campus, Stockton Road, Durham, DH1 3LE, United Kingdom • *E-mail* : katie.gittins@durham.ac.uk

CORENTIN LÉNA, Università degli Studi di Padova, Dipartimento di Tecnica e Gestione dei Sistemi Industriali (DTG), Stradella S. Nicola 3, 36100 Vicenza and Dipartimento di Matematica “Tullio Levi-Civita”, via Trieste 63, 35121 Padova, Italy • *E-mail* : corentin.lena@unipd.it

Mathematical subject classification (2010). — 35P15, 35J05, 49R05.

Key words and phrases. — Courant's nodal domain theorem, Neumann eigenvalues, Robin eigenvalues, Euclidean domains.

and not in \mathbb{R}^2 . It is not a priori clear that the Faber–Krahn inequality with the same constant as in the Euclidean case (hereafter called *classical Faber–Krahn inequality*) holds for V^R . The same gap occurs in Section 5.3 when considering a possibly multiply connected domain in \mathbb{R}^2 , for which V^R is an open set in the disjoint union $\bigcup_{i=0}^b \mathcal{C}_{L_i}$. Similarly, when $\Omega \subset \mathbb{R}^n$, we implicitly apply (on page 123) the classical Faber–Krahn inequality in the cylindrical manifold $N(\Gamma) = \Gamma \times \mathbb{R}$, where it might not hold.

When Ω is a (possibly multiply connected) C^2 domain in \mathbb{R}^2 , we can fix the issue by combining a few known geometric results. Examination of the proof of the classical Faber–Krahn inequality by symmetrization (see, for instance, [1, I.9]) shows that it holds provided that all open sets $W \subset V^R$ satisfy the classical isoperimetric inequality $|\partial W|^2 \geq 4\pi|W|$. From [3, §6], the classical isoperimetric inequality holds for any open set $W_i \subset \mathcal{C}_{L_i}$ such that $|W_i| \leq L_i^2/\pi$. Note that the parameter a in [3] is the radius of the cylinder, equal to $L_i/2\pi$ in our notation. We then remark that V^R can be written as a disjoint union $\bigcup_{i=0}^b V_i$, with each V_i contained in

$$(\mathbb{R}/(L_i\mathbb{Z})) \times (-\delta_0(\Omega), \delta_0(\Omega)) \subset \mathcal{C}_{L_i}$$

(some V_i 's may be empty). In addition, $\delta_0(\Omega)$ is less than or equal to the smallest radius of curvature of $\partial\Omega$, denoted by $t_+(\Omega)$. We can write $\Omega = \Omega_0 \setminus \bigcup_{i=1}^b \overline{D}_i$, where Ω_0 and all the D_i 's are simply connected C^2 domains and apply to each the inequality $\pi t_+(D)^2 \leq |D|$, valid for any simply connected domain $D \subset \mathbb{R}^2$ (see [4] and references therein). We obtain

$$\delta_0(\Omega) \leq t_+(\Omega) \leq \frac{1}{\pi^{1/2}} \min \left\{ |\Omega_0|^{1/2}, \min_{1 \leq i \leq b} |D_i|^{1/2} \right\} \leq \frac{1}{2\pi} \min_{0 \leq i \leq b} L_i$$

(where the last inequality follows from the isoperimetric inequality in \mathbb{R}^2). Thus, $|V_i| \leq 2\delta_0(\Omega)L_i \leq L_i^2/\pi$ for all i , and any $W_i \subset V_i$ satisfies the isoperimetric inequality. This is, therefore, also the case for any $W = \bigcup_{i=0}^b W_i \subset V^R$, which concludes the proof. All the results of Sections 2 to 8 hold without any change.

Most of the material in Section 9 can be recovered by replacing the quantities $\delta_+(x')$, $\underline{\delta}_+(\Omega)$ and $\delta_0(\Omega)$, defined on page 121, with, respectively,

$$\begin{aligned} \delta(x') &:= \sup\{\delta > 0 : \text{dist}(F(x', t), \partial\Omega) = |t| \text{ for all } t \in [-\delta, \delta]\}, \\ \underline{\delta}(\Omega) &:= \inf_{x' \in \partial\Omega} \delta(x'), \\ \delta_1(\Omega) &:= \min\{t_+(\Omega), \underline{\delta}(\Omega)\}. \end{aligned}$$

Having defined the set $V^R \subset \Gamma \times \mathbb{R}$ as in Section 5.2, we modify our method by taking the additional steps of considering the image $U^R := F(V^R) \subset \mathbb{R}^n$ and applying the Faber–Krahn inequality to U^R . We get an upper bound on the number of boundary nodal domains $\nu_1(\varepsilon_0, u)$, and thus find results similar

to Proposition 9.2, except that $\delta_0(\Omega)$ is replaced with $\delta_1(\Omega)$, and the constants depending only on n are larger ones. Note that the geometric quantity $\delta_1(\Omega)$ depends on the cut-distance to the boundary (also known as the injectivity radius of the normal exponential map) with respect to both the interior and the exterior of the domain Ω , whereas $\delta_0(\Omega)$ depends only on the interior. If, in addition, Ω is assumed to be convex, then $\delta_1(\Omega) = \delta_0(\Omega)$, since $\text{dist}(F(x', t), \partial\Omega) = -t$ for all $x' \in \partial\Omega$ and $t \leq 0$ (the cut-distance with respect to the exterior is infinite). Therefore, in that case, only the constants depending on n change. In particular, Proposition 9.4 holds as written.

BIBLIOGRAPHY

- [1] P. BÉRARD & D. MEYER – “Inégalités isopérimétriques et applications. (Isoperimetric inequalities and applications)”, *Ann. Sci. Éc. Norm. Supér. (4)* **15** (1982), p. 513–541.
- [2] K. GITTINS & C. LÉNA – “Upper bounds for Courant-sharp Neumann and Robin eigenvalues”, *Bull. Soc. Math. Fr.* **148** (2020), no. 1, p. 99–132.
- [3] H. HOWARDS, M. HUTCHINGS & F. MORGAN – “The isoperimetric problem on surfaces”, *Am. Math. Mon.* **106** (1999), no. 5, p. 430–439.
- [4] K. PANKRASHKIN – “An inequality for the maximum curvature through a geometric flow”, *Arch. Math.* **105** (2015), no. 3, p. 297–300.