

quatrième série - tome 55 fascicule 1 janvier-février 2022

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

YVES DE CORNULIER

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} octobre 2021

S. CANTAT G. GIACOMIN
G. CARRON D. HÄFNER
Y. CORNULIER D. HARARI
F. DÉGLISE C. IMBERT
A. DUCROS S. MOREL
B. FAYAD P. SHAN

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.

Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.

annaes@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France

Case 916 - Luminy

13288 Marseille Cedex 09

Tél. : (33) 04 91 26 74 64

Fax : (33) 04 91 41 17 51

email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 437 euros.

Abonnement avec supplément papier :

Europe : 600 €. Hors Europe : 686 € (\$ 985). Vente au numéro : 77 €.

© 2022 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Fabien Durand

Périodicité : 6 n^{os} / an

OPENNESS OF UNIFORM K-STABILITY IN FAMILIES OF \mathbb{Q} -FANO VARIETIES

BY HAROLD BLUM AND YUCHEN LIU

ABSTRACT. – We show that uniform K-stability is a Zariski open condition in \mathbb{Q} -Gorenstein families of \mathbb{Q} -Fano varieties. To prove this result, we consider the behavior of the stability threshold in families. The stability threshold (also known as the delta-invariant) is a recently introduced invariant that is known to detect the K-semistability and uniform K-stability of a \mathbb{Q} -Fano variety. We show that the stability threshold is lower semicontinuous in families and provides an interpretation of the invariant in terms of the K-stability of log pairs.

RÉSUMÉ. – Nous démontrons que la K-stabilité uniforme est une condition ouverte pour la topologie de Zariski dans une famille \mathbb{Q} -Gorenstein de variétés algébriques \mathbb{Q} -Fano. Pour prouver ce résultat, nous considérons le seuil de stabilité en familles. Le seuil de stabilité (également appelé l'invariant delta) est un invariant qui détecte la K-semistabilité et la K-stabilité uniforme d'une variété algébrique \mathbb{Q} -Fano. Nous démontrons que le seuil de stabilité est semi-continu inférieurement en familles et fournissons une interprétation de cet invariant en termes de K-stabilité des paires log.

Throughout, we work over a characteristic zero algebraically closed field k .

1. Introduction

In this article, we consider the behavior of K-stability in families of \mathbb{Q} -Fano varieties. Recall that K-stability is an algebraic notion introduced by Tian [53] and later reformulated by Donaldson [21] to detect the existence of certain canonical metrics on complex projective varieties. In the special case of complex \mathbb{Q} -Fano varieties, the Yau-Tian-Donaldson conjecture states that a complex \mathbb{Q} -Fano variety is K-polystable iff it admits a Kähler-Einstein metric. (By a \mathbb{Q} -Fano variety, we mean a projective variety that has at worst klt singularities and anti-ample canonical divisor.) For smooth complex Fano varieties, this conjecture was recently settled in the work of Chen-Donaldson-Sun and Tian [16, 54] (see also [19, 17, 2]).

Another motivation for understanding the K-stability of \mathbb{Q} -Fano varieties is to construct compact moduli spaces for such varieties. It is expected that there is a proper good moduli space parametrizing K-polystable \mathbb{Q} -Fano varieties of fixed dimension and volume. For

smoothable \mathbb{Q} -Fano varieties, such a moduli space is known to exist [40] (see also [48, 43]). A key step in constructing the moduli space of \mathbf{K} -polystable Fano varieties is verifying the Zariski openness of \mathbf{K} -semistability. Towards this goal, we prove

THEOREM A. – *If $\pi : X \rightarrow T$ is a projective flat family of varieties such that T is normal, π has normal connected fibers, and $-K_{X/T}$ is \mathbb{Q} -Cartier and π -ample, then*

- (1) $\{t \in T \mid X_{\bar{t}} \text{ is uniformly } \mathbf{K}\text{-stable}\}$ is a Zariski open subset of T , and
- (2) $\{t \in T \mid X_{\bar{t}} \text{ is } \mathbf{K}\text{-semistable}\}$ is a countable intersection of Zariski open subsets of T .

The notion of uniform \mathbf{K} -stability is a strengthening of \mathbf{K} -stability introduced in [11, 20]. In [2], it was shown that a smooth Fano variety X with discrete automorphism group is uniformly \mathbf{K} -stable iff there exists a Kähler-Einstein metric on X . The latter equivalence was later extended to \mathbb{Q} -Fano varieties with discrete automorphism group in [55]. \mathbf{K} -semistability is strictly weaker than \mathbf{K} -(poly)stability and corresponds to being almost Kähler-Einstein [37, 2].

In [7], the first author and Xu show that the moduli functor of uniformly \mathbf{K} -stable \mathbb{Q} -Fano varieties of fixed volume and dimension is represented by a separated Deligne-Mumford stack, which has a coarse moduli space that is a separated algebraic space. The proof of the result combines Theorem A.1 with a boundedness statement in [28] (that uses ideas from [3]) and a separatedness statement in [7].

For smooth families of Fano varieties, Theorem A is not new. Indeed, for a smooth family of complex Fano varieties with discrete automorphism group, the \mathbf{K} -stable locus is Zariski open by [44, 23]. In [40], it was shown that the \mathbf{K} -semistable locus is Zariski open in families of smoothable \mathbb{Q} -Fano varieties. These results all rely on deep analytic tools developed in [16, 54].

Unlike the previous results, our proof of Theorem A is purely algebraic. (A different algebraic proof of Theorem A.2 was also given in [6] using a characterization of \mathbf{K} -semistability in terms of the normalized volume of the affine cone over a \mathbb{Q} -Fano variety [36, 38, 41].) Furthermore, the result holds for all \mathbb{Q} -Fano varieties, including those that are not smooth(able), and also log Fano pairs. The argument relies on new tools for characterizing the uniform \mathbf{K} -stability and \mathbf{K} -semistability of Fano varieties [11, 36, 24, 25, 5].

More precisely, our approach to proving Theorem A is through understanding the behavior of the *stability threshold* (also known as δ -invariant or *basis log canonical threshold*) in families. We recall the definition of this new invariant.

Let X be a projective klt variety and L an ample Cartier divisor on X . Set

$$|L|_{\mathbb{Q}} := \{D \in \text{Div}(X)_{\mathbb{Q}} \mid D \geq 0 \text{ and } mD \sim mL \text{ for some } m \in \mathbb{Z}_{>0}\}.$$

Following [25], we say that $D \in |L|_{\mathbb{Q}}$ is an m -basis type divisor of L if there exists a basis $\{s_1, \dots, s_{N_m}\}$ of $H^0(X, \mathcal{O}_X(mL))$ such that

$$D = \frac{1}{mN_m} (\{s_1 = 0\} + \dots + \{s_{N_m} = 0\}).$$

For $m \in M(L) := \{m \mid h^0(X, \mathcal{O}_X(mL)) \neq 0\}$, set

$$\delta_m(X; L) := \inf_{D \text{ } m\text{-basis type}} \text{lct}(X; D),$$

where $\text{lct}(X; D)$ denotes the log canonical threshold of D . The stability threshold of L is

$$\delta(X; L) := \limsup_{M(L) \ni m \rightarrow \infty} \delta_m(X; L).$$

In fact, the above limsup is a limit by [5]. If X is a \mathbb{Q} -Fano variety, we set

$$\delta(X) := r\delta(X; -rK_X),$$

where $r \in \mathbb{Z}_{>0}$ is such that $-rK_X$ is Cartier. (The definition is independent of the choice of r .)

The stability threshold is closely related to the *global log canonical threshold of L* , which is an algebraic version of Tian's α -invariant. Recall that the global log canonical threshold of L is

$$\alpha(X; L) := \inf_{D \in |L|_{\mathbb{Q}}} \text{lct}(X; D).$$

The two thresholds satisfy

$$\frac{n+1}{n}\alpha(X; L) \leq \delta(X; L) \leq (n+1)\alpha(X; L),$$

where $n = \dim(X)$.

The stability threshold was introduced in the \mathbb{Q} -Fano case by K. Fujita and Y. Odaka to characterize the K-stability of \mathbb{Q} -Fano varieties [25]. More generally, the invariant coincides with an invariant suggested by R. Berman and defined in [12]. Using the valuative criterion for K-stability of K. Fujita and C. Li [24, 36], it was shown that the invariant characterizes certain K-stability notions.

THEOREM 1.1 ([25, 5]). – *Let X be a \mathbb{Q} -Fano variety.*

- (1) *X is uniformly K-stable iff $\delta(X) > 1$.*
- (2) *X is K-semistable iff $\delta(X) \geq 1$.*

In light of the previous statement, Theorem A is a consequence of the following result.

THEOREM B. – *Let $\pi : X \rightarrow T$ be a projective flat family of varieties and L a π -ample Cartier divisor on X . Assume T is normal, X_t is a klt variety for all $t \in T$, and $K_{X/T}$ is \mathbb{Q} -Cartier. Then, the two functions $T \rightarrow \mathbb{R}$ defined by*

$$T \ni t \mapsto \delta(X_{\bar{t}}; L_{\bar{t}}) \quad \text{and} \quad T \ni t \mapsto \alpha(X_{\bar{t}}; L_{\bar{t}})$$

are lower semicontinuous.

In the above theorem, $(X_{\bar{t}}; L_{\bar{t}})$ denotes the restriction of (X, L) to the geometric fiber over t . As explained in [18, Remark 4.15], the above result would not hold with “ $\delta(X_{\bar{t}}; L_{\bar{t}})$ ” replaced by “ $\delta(X_t; L_t)$.”

Let us note the main limitation of Theorem B. While the statement implies that $\{t \in T \mid \delta(X_{\bar{t}}; L_{\bar{t}}) > a\}$ is open for each $a \in \mathbb{R}_{\geq 0}$, it does not imply the fact that $t \mapsto \delta(X_{\bar{t}}; L_{\bar{t}})$ takes finitely many values. Hence, we are unable to prove $\{t \in T \mid \delta(X_{\bar{t}}; L_{\bar{t}}) \geq a\}$ is open and cannot verify the openness of K-semistability in families of \mathbb{Q} -Fano varieties. The openness of K-semistability is an immediate consequence of Theorem B and the following conjecture (see [6, Conjecture 2] for a local analogue).