

ON SEMI-STABLE, SINGULAR CUBIC SURFACES

by

Nguyen Chanh Tu

Abstract. — This paper deals with semi-stable and stable singular cubic surfaces from the point of view of the geometric invariant theory. We are interested in properties of the subsets iA_1jA_2 corresponding to all semi-stable, singular cubic surfaces with exactly i singular points of type A_1 and j singular points of type A_2 . We consider semi-stable cubic surfaces as “c-surfaces” of 6-point schemes in almost general position with some conditions of configurations. This is a generalization of the blowing-up of $\mathbb{P}^2$ at 6 points in general position. From relevant configurations of 6-point schemes, we can determine number of star points, the configuration of singular points, of lines and tritangent planes with multiplicities on semi-stable, singular cubic surfaces.

Résumé (Sur les surfaces cubiques semi-stables). — Cet article concerne les surfaces cubiques semi-stables et stables du point de vue de la théorie géométrique des invariants. Nous nous sommes intéressés aux propriétés des sous-ensembles iA_1jA_2 correspondant à toutes les surfaces cubiques singulières semi-stables avec exactement i points singuliers de type A_1 et j points singuliers de type A_2 . Nous considérons les surfaces cubiques semi-stables comme « c-surfaces » d'ensembles de 6 points en position presque générale avec certaines conditions de configurations. Ceci est une généralisation de l'éclatement de $\mathbb{P}^2$ en 6 points en position générale. À partir de configurations adaptées d'ensembles de 6 points, nous pouvons déterminer le nombre de points « étoile », la configuration des points singuliers, des droites et des plans « tritangents » avec multiplicités sur les surfaces singulières cubiques semi-stables.

1. Introduction

Consider $\mathbb{P}^{19}$ as a parametrizing space of cubic surfaces in $\mathbb{P}_k^3$, where k is an algebraically closed field with characteristic 0. We have the action of $\mathrm{PGL}(4)$ on $\mathbb{P}^{19}$. The locus $\Delta \subset \mathbb{P}^{19}$ of singular cubic surfaces is a closed subset of codimension 1. Some

2000 Mathematics Subject Classification. — Primary 14C05, 14J17, 14J10; Secondary 14C20, 14J25.

Key words and phrases. — Varieties and morphism, special surfaces, singularity.

The author is supported by Japan Society for the Promotion of Science (JSPS) Postdoctoral Fellowship and by Ministry of Education and Training of Vietnam.

classifications of singular cubic surfaces can be found in [4] or [5]. We are interested in singular cubic surfaces which correspond to semi-stable and stable points under the action of $\mathrm{PGL}(4)$ on $\mathbb{P}^{19}$ in the sense of the geometric invariant theory. One reason we are interested in these kinds of singularities is that the quotient space of semi-stable points over $\mathrm{PGL}(4)$ exists and it is a compactification of the moduli space of non-singular cubic surfaces.

It is well-known that the blowing-up of $\mathbb{P}^2$ at 6 points in general position is isomorphic to a non-singular cubic surface. Conversely, any non-singular cubic surface can be obtained in that way. A question arises naturally: is there a similar correspondence between a semi-stable, singular cubic surface and a 6-point scheme in some relevant configuration of its points? Showing such a correspondence is one of main goals of this paper. Namely, let X be a semi-stable cubic surface. Then there exists a 6-point scheme $\mathcal{P}$ such that the linear system $\mathcal{L}_{\mathcal{P}}$ of cubic forms in four variables through $\mathcal{P}$ has dimension 4; furthermore, for any basis of $\mathcal{L}_{\mathcal{P}}$, the closure of the image of the rational map from $\mathbb{P}^2$ to $\mathbb{P}^3$ defined by the basis is a surface which is isomorphic to X . In this case, we have a morphism $Y \rightarrow X$, where Y is the blowing-up of $\mathbb{P}^2$ at $\mathcal{P}$. In general, this is a blowing-down and not an isomorphism. A close study of such 6-point schemes enables us to determine the number of lines, the number of singularities of X and their configuration as well.

This also gives a way to compute the multiplicity of lines and tritangent planes on semi-stable, singular cubic surfaces. This investigation shows a clear picture on the configuration of lines and tritangent planes of semi-stable, singular cubic surfaces. Moreover, we will give definitions of *star point* and *proper star point* which are generalizations of the concept of Eckardt point on non-singular cubic surfaces. We will determine the number of (proper) star points on a general one of any class of semi-stable cubic surfaces and study some properties.

2. Stable and semi-stable, singular cubic surfaces

We denote by iA_1jA_2 the subset of $\mathbb{P}^{19}$ corresponding to irreducible cubic surfaces with exactly i singular points of type A_1 and j singular points of type A_2 . We refer to [1] and [2] or to [4] for general definitions of types of singularities. We will see later that these subsets correspond to all semi-stable, singular cubic surfaces with respect to the action of $\mathrm{PGL}(4)$ on $\mathbb{P}^{19}$.

Remark 2.1

(i) In the case of cubic surfaces, the singularities of types A_1 and A_2 are characterized as follows. A point P on a cubic surface with only isolated singularities is called a *singular point of type A_1* (respectively A_2) if the tangent cone at P is an irreducible quadric surface (respectively if the tangent cone at P consists of two distinct planes whose intersection line does not lie on the surface).

(ii) We have $2i + 3j \leq 9$, $i \leq 4$ and $(i, j) \neq (3, 1)$, see [4], p. 255 or [11], pp. 49-50. We use $j\mathcal{A}_2$ and $i\mathcal{A}_1$ instead of $0\mathcal{A}_1j\mathcal{A}_2$ and $i\mathcal{A}_10\mathcal{A}_2$, respectively.

(iii) By the definition, a semi-stable, singular cubic surface can be given by a polynomial in the following form:

$$F = x_3f_2(x_0, x_1, x_2) + f_3(x_0, x_1, x_2),$$

where f_i for $i = 1, 2$ is a homogeneous polynomial of degree i . Then the type of singularity of the surface is characterized by $\text{rank}(f_2)$ and the configuration of points in $V_{\mathbb{P}^2}(f_2, f_3)$.

Some interesting properties of subsets $i\mathcal{A}_1j\mathcal{A}_2$ are shown in the following.

Proposition 2.2. — *The subsets $i\mathcal{A}_1j\mathcal{A}_2$ are irreducible of codimension $i + 2j$ in $\mathbb{P}^{19}$ and have a relation as shown in the Figure 1, where $A \longrightarrow B$ means that $\overline{A} \subset \overline{B}$ and subsets are in the same column iff they have the same codimension.*




FIGURE 1

Proof. — This follows from [3], Prop. 2.1. and [3], Fig. 1, p. 435. $\square$

Proposition 2.3. — *On the action of $\text{PGL}(4)$ on $\mathbb{P}^{19}$, we have:*

- (i) *The subset of stable points consists of points in $\mathbb{P}^{19} - \Delta$ and those of types $i\mathcal{A}_1$ for $1 \leq i \leq 4$.*
- (ii) *The subset of semi-stable points consists of points in $\mathbb{P}^{19} - \Delta$ and all those of types $i\mathcal{A}_1j\mathcal{A}_2$.*

Proof. — This result was mentioned, for instance, in [10], p. 80 or [9], p. 51. A detailed proof could be found in [11], 3.2.14. $\square$

3. Semi-stable as csurfaces of 6-point schemes in almost general position

As in the case of non-singular cubic surfaces, we show that each semi-stable, singular cubic surface corresponds to a relevant 6-point scheme in almost general position. Moreover we prove that the corresponding semi-stable cubic surfaces are isomorphic if their 6-points schemes are different by quadratic transformations.

Definition. — A 6-point scheme is a closed subscheme in $\mathbb{P}^2$ of dimension zero and of length 6. Any 6-point scheme $\mathcal{P}$ defines a formal cycle $c(\mathcal{P}) = \sum n_i P_i$ for $\sum n_i = 6$; the set of the points P_i is called *the support* of $\mathcal{P}$ and denoted by $\text{Supp}(\mathcal{P})$. If the linear system of all cubic forms passing through a 6-point scheme $\mathcal{P}$ has (linear) dimension 4, then $\mathcal{P}$ is called a 6-point scheme *in almost general position*.

Let Hilb_n denote the Hilbert scheme of zero-dimensional closed subschemes of length n in $\mathbb{P}^2$. We denote by $\mathcal{H}^a$ the subscheme of Hilb_6 consisting of all 6-point schemes in almost general position.

Let $\mathcal{P} \in \mathcal{H}^a$ and let l be any line in $\mathbb{P}^2$ such that $l \cap \mathcal{P} \neq \emptyset$. Then the length of $l \cap \mathcal{P}$ is not greater than 4.

Definition. — Let $\mathcal{P} \in \mathcal{H}^a$. We say that $\mathcal{P}$ is a 6-point scheme with *no 4 points on a line* if there does not exist any line l in $\mathbb{P}^2$ such that the length of $l \cap \mathcal{P}$ is equal to 4. Denote by $\mathcal{H}^o$ the subset of 6-point schemes with no 4 points on a line.

Lemma 3.1. — Let $\mathcal{P} \in \mathcal{H}^o$. Let $\mathcal{L}_{\mathcal{P}}$ be the linear system of cubic forms passing through $\mathcal{P}$.

- (i) The base locus of $\mathcal{L}_{\mathcal{P}}$ is the support of $\mathcal{P}$.
- (ii) Let $\{f_1, \dots, f_4\}$ be a basis of $\mathcal{L}_{\mathcal{P}}$. Consider the morphism

$$\begin{aligned} \psi : \mathbb{P}^2 - \text{Supp}(\mathcal{P}) &\longrightarrow \mathbb{P}^3 \\ P &\longmapsto (f_1(P) : f_2(P) : f_3(P) : f_4(P)). \end{aligned}$$

Let X be the closure of the image of ψ . Then X is a cubic surface.

- (iii) If $\{g_1, \dots, g_4\}$ is another basis of $\mathcal{L}_{\mathcal{P}}$ and X' is the cubic surface obtained as in (ii), then X and X' are isomorphic.

Proof

(i) Let $P \in \mathbb{P}^2 - \text{Supp}(\mathcal{P})$. Since $\mathcal{P}$ does not have 4 points on a line, there exists a cubic form in $\mathcal{L}_{\mathcal{P}}$ which does not contain P . This implies that the base locus of $\mathcal{L}_{\mathcal{P}}$ is the support of $\mathcal{P}$.

(ii) Let Q_1, Q_2 be two general points in $\mathbb{P}^2 - \text{Supp}(\mathcal{P})$. The linear subspaces consisting of cubic forms through $\mathcal{P} \cup \{Q_1\}$ and $\mathcal{P} \cup \{Q_1, Q_2\}$ respectively have dimension 3 and 2. This implies that there exists a cubic form in $\mathcal{L}_{\mathcal{P}}$ which contains Q_1 but does not contain Q_2 and conversely. This means that ψ is injective over an open subset of $\mathbb{P}^2$. Moreover, any two general cubic forms in $\mathcal{L}_{\mathcal{P}}$ have 3 other points in common which do not belong to $\mathcal{P}$. This implies that X is a cubic surface.

(iii) Let $A = (a_{ij})_{4 \times 4}$ be the base change matrix from $\{f_1, \dots, f_4\}$ to $\{g_1, \dots, g_4\}$. Then A defines a projective transformation which transforms X to X' . $\square$

Definition. — A *csurface* is an algebraic variety Y such that there exists a cubic surface $X \subset \mathbb{P}^3$ such that $X \cong Y$. From the lemma, we see that each $\mathcal{P} \in \mathcal{H}^o$ determines uniquely (up to isomorphisms) a csurface, which is called *the csurface*

of $\mathcal{P}$. If $\mathcal{P}$ consists of 6 points in general position, then the csurface of $\mathcal{P}$ is the blowing-up of $\mathbb{P}^2$ at $\mathcal{P}$.

Definition. — Let $P_0 = (1 : 0 : 0)$, $P_1 = (0 : 1 : 0)$ and $P_2 = (0 : 0 : 1)$. Let $\varphi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ be the quadratic transformation with respect to P_0, P_1 and P_2 (see [8], V.4.2.3). Let C be the cubic curve given by

$$(1) \quad F = \sum a_{ijk} x_0^i x_1^j x_2^k \text{ for } i+j+k=3 \text{ and } 0 \leq i, j, k \leq 2.$$

The cubic curve defined by $F_\varphi := \sum a_{ijk} y_0^{2-i} y_1^{2-j} y_2^{2-k}$ in $\mathbb{P}^2$ is called *the image of C by φ* and is denoted by C_φ .

Lemma 3.2. — Let $\mathcal{P} \in \mathcal{H}^o$. Suppose that $\text{Supp}(\mathcal{P})$ contains 3 distinct points P_1, P_2 and P_3 . Suppose further that there exists a cubic form in $\mathcal{L}_{\mathcal{P}}$ which is non-singular at any P_i for $i = 1, 2, 3$. Let φ be the quadratic transformation with respect to P_1, P_2 and P_3 . Then the set $\varphi(\mathcal{L}_{\mathcal{P}}) := \{F_\varphi \mid F \in \mathcal{L}_{\mathcal{P}}\}$ is a 4-dimensional linear space whose base locus is of dimension 0.

Proof. — Choose coordinates such that $P_1 = (1 : 0 : 0)$, $P_2 = (0 : 1 : 0)$ and $P_3 = (0 : 0 : 1)$. Suppose that the base locus of $\varphi(\mathcal{L}_{\mathcal{P}})$ contains an irreducible component Y of positive dimension. Since φ is one-to-one in $\mathbb{P}^2 - V(x_0 x_1 x_2)$, the variety Y is contained in $V(y_0 y_1 y_2)$. Assume that Y contains the line $d_{12} = V(y_0)$. This means that for any $F \in \mathcal{L}_{\mathcal{P}}$, we have $F_\varphi = y_0 g_2(y_0, y_1, y_2)$ where g_2 is a homogeneous polynomial of degree 2 and vanishes at $Q_3 = (0 : 0 : 1)$. Then $F = (F_\varphi)_{\varphi^{-1}}$ is singular at $P_1 = (1 : 0 : 0)$. A contradiction! $\square$

Definition. — Let $\mathcal{P} \in \mathcal{H}^o$ satisfy the conditions as in the previous lemma. Let I be the ideal generated by all cubic forms in $\varphi(\mathcal{L}_{\mathcal{P}})$. The scheme defined by this ideal is called *the image of $\mathcal{P}$* and denoted by $\varphi(\mathcal{P})$.

Proposition 3.3. — Every semi-stable cubic surface is isomorphic to the csurface of some 6-point scheme in almost general position with no 4 points on a line.

Proof. — Let X be a semi-stable cubic surface. If X is a non-singular cubic surface then it is isomorphic to the blowing-up of a 6-point scheme in general position. We consider the case that X is singular.

Suppose that X does not have any A_2 singularity. By choosing coordinates, we may assume X to be defined by

$$F = x_3 f_2(x_0, x_1, x_2) + f_3(x_0, x_1, x_2),$$

where f_i for $i = 2, 3$ is a homogeneous polynomial of degree i and f_2 is irreducible. The scheme $\mathcal{P} = V_{\mathbb{P}^2}(f_2, f_3)$ defines an element in $\mathcal{H}^o$. The 6-point scheme $\mathcal{P}$ is contained in an irreducible conic curve defined by f_2 and the cycle $c(\mathcal{P})$ corresponds to a partition $(2^{i-1}1^k)$ of 6. Let $\mathcal{L}_{\mathcal{P}}$ be the linear space of cubic forms passing through $\mathcal{P}$. Since $\mathcal{P}$ does not contain any triple point, we see that the cubic forms