

quatrième série - tome 51 fascicule 5 septembre-octobre 2018

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Jan NEKOVÁŘ

*Eichler-Shimura relations and semisimplicity
of étale cohomology of quaternionic Shimura varieties*

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Patrick BERNARD

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE

de 1883 à 1888 par H. DEBRAY

de 1889 à 1900 par C. HERMITE

de 1901 à 1917 par G. DARBOUX

de 1918 à 1941 par É. PICARD

de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} mars 2018

P. BERNARD

A. NEVES

S. BOUCKSOM

J. SZEFTTEL

R. CERF

S. VŨ NGỌC

G. CHENEVIER

A. WIENHARD

Y. DE CORNULIER G. WILLIAMSON

E. KOWALSKI

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,

45, rue d'Ulm, 75230 Paris Cedex 05, France.

Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.

annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France

Case 916 - Luminy

13288 Marseille Cedex 09

Tél. : (33) 04 91 26 74 64

Fax : (33) 04 91 41 17 51

email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 420 euros.

Abonnement avec supplément papier :

Europe : 540 €. Hors Europe : 595 € (\$ 863). Vente au numéro : 77 €.

© 2018 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Stéphane Seuret

Périodicité : 6 n^{os} / an

EICHLER-SHIMURA RELATIONS AND SEMISIMPLICITY OF ÉTALE COHOMOLOGY OF QUATERNIONIC SHIMURA VARIETIES

BY JAN NEKOVÁŘ

ABSTRACT. – We show that the non CM part of ℓ -adic étale cohomology of any compact quaternionic Shimura variety with coefficients in any automorphic local system is a semisimple Galois representation. If the local system has weight $k = (k_1, \dots, k_d)$ with all k_i of the same parity, the full ℓ -adic étale cohomology is semisimple. For Hilbert modular varieties, analogous results are proved for ℓ -adic intersection cohomology of the Baily-Borel compactification. The proof combines a representation-theoretical criterion of semisimplicity with Eichler-Shimura relations for partial Frobenius morphisms.

RÉSUMÉ. – On montre que l'action galoisienne sur la partie sans multiplication complexe de la cohomologie étale d'un faisceau ℓ -adique lisse automorphe sur une variété de Shimura quaternionique compacte est semi-simple. Si le poids du faisceau s'écrit $k = (k_1, \dots, k_d)$, où les k_i ont la même parité, toute la cohomologie étale est semi-simple. Les mêmes résultats sont montrés pour la cohomologie d'intersection ℓ -adique de la compactification de Baily-Borel des variétés modulaires de Hilbert. La preuve utilise un critère abstrait de semi-simplicité et les relations d'Eichler-Shimura pour les morphismes de Frobenius partiels.

0. Introduction

0.1. General conventions and notation

The characteristic polynomial of an endomorphism u of a finite-dimensional vector space over a field k will be denoted by $P_u(X) = \det(X \cdot \text{id} - u) \in k[X]$. If $k \subset K$ are fields and X is a k -vector subspace of a K -vector space Y , we denote by $K \cdot X$ the K -vector subspace of Y generated by X . We abbreviate $\otimes_{\mathbb{Z}}$ as $\otimes$. For an abelian group A we let $\widehat{A} = A \otimes \widehat{\mathbb{Z}}$. We denote by $\mathbf{A}$ and $\mathbf{A}_k$, respectively, the ring of adèles of $\mathbb{Q}$ and of a number field k .

Throughout the article we fix an isomorphism $\mathbb{C} \xrightarrow{\sim} \overline{\mathbb{Q}}_{\ell}$. For any algebraic object $(-)$ defined over a subfield of $\mathbb{C}$ we denote by $(-)_{\ell}$ its base change to $\overline{\mathbb{Q}}_{\ell}$. Let $\overline{\mathbb{Q}}$ be the algebraic closure of $\mathbb{Q}$ in $\mathbb{C}$. The reciprocity map of class field theory is normalized by letting uniformisers correspond to geometric Frobenius elements $\text{Fr}(P)$. All representations and

The author's research was supported in part by the grant ANR-BLAN-0114.

characters are assumed to be continuous with respect to the natural topologies involved. A representation of a profinite group is called strongly irreducible if its restriction to every open subgroup is irreducible. By an automorphic representation we mean an irreducible automorphic representation.

0.2. – Let us recall basic facts about decomposition of singular and étale cohomology of compact Shimura varieties. As in the classical case of cuspidal cohomology of modular curves, everything boils down to the fact that the Hecke operators act on the space of cuspidal automorphic forms in a selfadjoint way (up to a twist).

Let $(G, \mathcal{X})$ be a (pure) Shimura datum. A rational representation $\xi : G_{\mathbb{C}} \rightarrow GL(N)_{\mathbb{C}}$ (whose restriction to the center $Z_{\mathbb{C}}$ satisfies an appropriate condition) gives rise, for each sufficiently small open compact subgroup $K \subset G(\widehat{\mathbf{Q}})$, to a locally constant sheaf of complex vector spaces $\mathcal{L}_{\xi}$ on the complex manifold $\mathrm{Sh}_K(G, \mathcal{X})^{\mathrm{an}} = G(\mathbf{Q}) \backslash (\mathcal{X} \times G(\widehat{\mathbf{Q}})/K)$.

0.3. – If, in addition, the derived group G^{der} is anisotropic, then $\mathrm{Sh}_K(G, \mathcal{X})^{\mathrm{an}}$ is compact and its cohomology $H^*(\mathrm{Sh}_K(G, \mathcal{X})^{\mathrm{an}}, \mathcal{L}_{\xi})$ is described in terms of relative Lie algebra cohomology.

Write $\mathcal{X} = G(\mathbf{R})/K_{\infty}$, where K_{∞} is the stabilizer of a fixed base point in $\mathcal{X}$, and denote, for any $G(\mathbf{R})$ -module V , by V_0 the subspace of K_{∞} -finite vectors in V . There is a canonical isomorphism

$$\begin{aligned} H^i(\mathrm{Sh}_K(G, \mathcal{X})^{\mathrm{an}}, \mathcal{L}_{\xi}) \\ = H^i(\mathfrak{g}, K_{\infty}; C^{\infty}(G(\mathbf{Q}) \backslash G(\mathbf{A})/K) \otimes \xi) = H^i(\mathfrak{g}, K_{\infty}; C^{\infty}(G(\mathbf{Q}) \backslash G(\mathbf{A})/K)_0 \otimes \xi), \end{aligned}$$

where $\mathfrak{g} = \mathrm{Lie}(G(\mathbf{R}))$ [4, VII.2.7]. It gives rise to a $G(\widehat{\mathbf{Q}})$ -equivariant isomorphism (0.3.1)

$$H^i(\mathrm{Sh}(G, \mathcal{X})^{\mathrm{an}}, \mathcal{L}_{\xi}) = \varinjlim_K H^i(\mathrm{Sh}_K(G, \mathcal{X})^{\mathrm{an}}, \mathcal{L}_{\xi}) = H^i(\mathfrak{g}, K_{\infty}; C^{\infty}(G(\mathbf{Q}) \backslash G(\mathbf{A}))_0 \otimes \xi).$$

For every character $\omega : Z(\mathbf{Q}) \backslash Z(\mathbf{A}) \rightarrow \mathbf{C}^{\times}$ fix a character $\omega' : G(\mathbf{Q}) \backslash G(\mathbf{A}) \rightarrow \mathbf{R}_+^{\times}$ such that $\omega'|_{Z(\mathbf{A})} = |\omega|$. The space $G(\mathbf{Q}) \backslash G(\mathbf{A})$ is compact (since G^{der} is anisotropic) and the completion $L^2(G, \omega)$ of

$$C^{\infty}(G, \omega) = \{f \in C^{\infty}(G(\mathbf{Q}) \backslash G(\mathbf{A})) \mid f(gz) = \omega(z)f(g) \quad \forall z \in Z(\mathbf{A})\}$$

with respect to the norm

$$\|f\|^2 = \int_{G(\mathbf{Q}) \backslash Z(\mathbf{A}) \backslash G(\mathbf{A})} (\omega'(g)^{-1} |f(g)|)^2 dg$$

is a unitary representation of $G(\mathbf{A})$ under the action $(g * f)(h) = \omega'(g)^{-1} f(hg)$. This representation decomposes as a discrete Hilbert sum $L^2(G, \omega) = \widehat{\bigoplus} m(\pi') \pi'$ of unitary automorphic representations π' of $G(\mathbf{A})$ with finite multiplicities $m(\pi')$.

Each π' has central character $\omega_{\pi'} = \omega/|\omega|$ and gives rise to an automorphic representation $\pi = \omega' \pi' = \pi_{\infty} \otimes \pi^{\infty}$ of $G(\mathbf{A}) = G(\mathbf{R}) \times G(\widehat{\mathbf{Q}})$ with central character $\omega_{\pi} = \omega$.

Matsushima's formula [4, Thm. VII.5.2] yields a $G(\widehat{\mathbf{Q}})$ -equivariant isomorphism

$$H^i(\mathfrak{g}, K_{\infty}; C^{\infty}(G, \omega)_0 \otimes \xi) = \bigoplus_{\pi = \pi_{\infty} \otimes \pi^{\infty}} m(\pi) H^i(\mathfrak{g}, K_{\infty}; \pi_{\infty} \otimes \xi) \otimes \pi^{\infty},$$