

*quatrième série - tome 55    fascicule 5    septembre-octobre 2022*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Alexis MICHELAT & Tristan RIVIÈRE

*The Classification of Branched Willmore Spheres  
in the 3-Sphere and the 4-Sphere*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

YVES DE CORNULIER

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1<sup>er</sup> octobre 2021

S. CANTAT      G. GIACOMIN  
G. CARRON     D. HÄFNER  
Y. CORNULIER D. HARARI  
F. DÉGLISE    C. IMBERT  
A. DUCROS     S. MOREL  
B. FAYAD      P. SHAN

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
Email : [annales@ens.fr](mailto:annales@ens.fr)

---

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Tél. : (33) 04 91 26 74 64. Fax : (33) 04 91 41 17 51  
Email : [abonnements@smf.emath.fr](mailto:abonnements@smf.emath.fr)

Tarifs

Abonnement électronique : 441 euros.  
Abonnement avec supplément papier :  
Europe : 619 €. Hors Europe : 698 € (\$ 985). Vente au numéro : 77 €.

---

© 2022 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Fabien Durand  
Périodicité : 6 n<sup>os</sup> / an

# THE CLASSIFICATION OF BRANCHED WILLMORE SPHERES IN THE 3-SPHERE AND THE 4-SPHERE

BY ALEXIS MICHELAT AND TRISTAN RIVIÈRE

---

**ABSTRACT.** – We extend the classification of Robert Bryant of Willmore spheres in  $S^3$  to *variational* branched Willmore spheres  $S^3$  and show that they are inverse stereographic projections of complete minimal surfaces with finite total curvature in  $\mathbb{R}^3$  and vanishing flux. We also obtain a classification of *variational* branched Willmore spheres in  $S^4$ , generalizing a theorem of Sébastien Montiel. The other main result of the article is a new local criterion which implies that branched Willmore spheres are conformally minimal.

**RÉSUMÉ.** – Nous étendons la classification de Robert Bryant des sphères de Willmore de  $S^3$  aux sphères de Willmore branchées et *variationnelles* de  $S^3$  et montrons que ces dernières sont des projections stéréographiques inverses de surfaces minimales de  $\mathbb{R}^3$  ayant une courbure totale finie et un flux nul. Nous obtenons également une classification des sphères de Willmore branchées et *variationnelles* de  $S^4$ , généralisant ainsi un théorème de Sébastien Montiel. L'autre théorème principal de l'article est un critère local impliquant qu'une sphère de Willmore branchée est conformément minimale.

## 1. Introduction

### 1.1. Willmore functional and quantization of energy

This article primarily addresses the generalization of Bryant's classification of smooth Willmore immersions of the sphere  $S^2$  into  $S^3$  to branched immersions. Let  $\Sigma$  be a closed Riemann surface, and  $n \geq 3$  be a fixed integer. The Willmore energy on a smooth Riemannian manifold  $(M^n, h)$  with sectional curvature  $\tilde{K}_h$  is defined on any smooth immersion  $\vec{\Phi} : \Sigma \rightarrow M^n$  by

$$W_{M^n}(\vec{\Phi}) = \int_{\Sigma} (|\vec{H}_g|^2 + \tilde{K}_h) d\text{vol}_g,$$

where  $g = \vec{\Phi}^*h$  is the pull-back metric of  $(M^n, h)$  by  $\vec{\Phi}$ , and  $\vec{H}_{\vec{\Phi}}$  is the mean-curvature, that is the half-trace of the second fundamental form  $\vec{\mathbb{I}}$  the immersion, given by

$$\vec{H}_g = \frac{1}{2} \sum_{i,j=1}^2 g^{i,j} \vec{\mathbb{I}}_{i,j}.$$

The most basic property of the Willmore energy is its conformal invariance which can be stated as follows. For all conformal diffeomorphism  $\varphi : (M^n, h) \rightarrow (\widetilde{M}^n, \widetilde{h})$ , we have

$$W_{\widetilde{M}^n}(\varphi \circ \vec{\Phi}) = W_{M^n}(\vec{\Phi}).$$

However, in the special case of  $\mathbb{R}^n$ , if  $\iota_a : \mathbb{R}^n \setminus \{a\} \rightarrow \mathbb{R}^n \setminus \{a\}$  is the inversion centered at  $a \in \vec{\Phi}(\Sigma)$ , we do not have in general

$$W_{\mathbb{R}^n}(\iota_a \circ \vec{\Phi}) = W_{\mathbb{R}^n}(\vec{\Phi}),$$

while we have the equality for inversions with center outside of  $\vec{\Phi}(\Sigma)$ . Nevertheless, the quantity

$$\mathcal{W}(\vec{\Phi}) = \int_{\Sigma} (|\vec{H}_g|^2 - K_g) d\text{vol}_g,$$

where  $K_g$  is the intrinsic Gauss curvature of  $\vec{\Phi}$ , is invariant under every conformal transformation. Indeed, the 2-form

$$(|\vec{H}_g|^2 - K_g) d\text{vol}_g = |\vec{h}_0|_{WP}^2 d\text{vol}_g,$$

where  $\vec{h}_0$  is the Weingarten tensor and  $|\cdot|_{WP}$  designs the Weil-Petersson metric, is a *pointwise* invariant (see for example 7.3.1 [35]). We shall come back to this point when we will state Noether’s theorem for the Willmore energy (see for Example (89) in the proof of Theorem 3.8).

We now come to the critical points of the Willmore energy. Classically, they are the smooth immersions satisfying to the equation

$$(1) \quad \Delta_g^N \vec{H} - 2|\vec{H}|^2 \vec{H} + \mathcal{A}(\vec{H}) + \mathcal{R}(\vec{H}) = 0,$$

where  $\Delta_g^N$  is the Laplacian on the normal bundle,  $\mathcal{A}$  the Simons operator and  $\mathcal{R}$  a curvature operator, given by

$$\mathcal{A}(\vec{H}) = \sum_{i,j=1}^2 \langle \vec{\mathbb{I}}(\vec{\varepsilon}_i, \vec{\varepsilon}_j), \vec{H} \rangle \vec{\mathbb{I}}(\vec{\varepsilon}_i, \vec{\varepsilon}_j), \quad \mathcal{R}(\vec{H}) = \left( \sum_{i=1}^2 R(\vec{H}, \vec{\varepsilon}_i) \vec{\varepsilon}_i \right)^N,$$

where  $(\vec{\varepsilon}_1, \vec{\varepsilon}_2)$  is any local orthonormal moving frame, and  $R$  is the Riemann curvature tensor of  $(M^n, h)$ . However, for the natural regularity  $\vec{\Phi} \in W^{2,2}(\Sigma, M^n)$  this equation does not even have a distributional meaning, as it would require  $\vec{H} \in L^3_{\text{loc}}(\Sigma, TM^n)$ . The weakest possible setting to work with is the space of *weak immersions* (introduced in [30], [31])

$$\mathcal{E}(\Sigma, M^n) = W^{2,2} \cap W^{1,\infty}(\Sigma, M^n) \cap \left\{ \begin{array}{l} \vec{\Phi} : d\vec{\Phi}(x) \text{ has rank 2 for almost all } x \in \Sigma \\ \text{and } \inf_{\Sigma} |d\vec{\Phi} \wedge d\vec{\Phi}|_{g_0} > 0 \end{array} \right\}$$

for any fixed Riemannian metric  $g_0$  on  $\Sigma$ . In the rest of the introduction, we suppose that  $M^n = \mathbb{R}^n$  and that  $h$  is the standard flat Euclidean metric. The second author showed in the Willmore equation can be written in a conservative weak formulation.

THEOREM ([30], Theorem I.1 p. 4). – *Let  $\Sigma$  be a closed Riemann surface, and  $\vec{\Phi} : \Sigma \rightarrow \mathbb{R}^n$  be a smooth immersion. Then, (identifying 2-vectors and functions on  $\Sigma$ )*

$$(2) \quad \Delta_g^N \vec{H}_g - 2|\vec{H}_g|^2 \vec{H}_g + \mathcal{A}(\vec{H}_g) = d \left( *_g d \vec{H}_g - 3 *_g (d \vec{H}_g)^N + \star(\vec{H}_g \wedge d\vec{n}) \right),$$

where  $\vec{H}_g$  is the mean curvature of  $\vec{\Phi}$ , where  $*_g$  is the Hodge star operator on  $\Sigma$  for the metric  $g$ , and  $\star$  the Hodge star operator on  $\mathbb{R}^n$  for the flat metric.

As the 1-form under the exterior derivative in (2) is in  $W^{-1,2} + L^1$ , the right-hand side is well-defined in a distributional sense as an element of  $\mathcal{D}'(\Sigma)$ . If the left-hand side is not defined in general for  $\vec{\Phi} \in \mathcal{E}(\Sigma, \mathbb{R}^n)$ , this comes from the fact that to write it, one has to make a projection on the normal bundle, while the normal is only in  $W^{1,2}(\Sigma, \mathcal{G}_{n-2}(\mathbb{R}^n))$ , where  $\mathcal{G}_{n-2}(\mathbb{R}^n)$  denotes the oriented Grassmannian of  $(n - 2)$ -plans in  $\mathbb{R}^n$ . Computing the Euler-Lagrange equation for arbitrary variations allows one to recover the weak formulation of the right-hand side (see [24]). Furthermore, as we shall see, the conservative form of the Euler-Lagrange equation of Willmore energy is a consequence of Noether’s theorem (this last fact already appears in a paper by Yann Bernard ([1] Theorem I.2 p. 220)).

Furthermore, writing the Willmore equation as the closeness of a 1-form allows one to introduce the concept of *variational* Willmore surfaces. In general, a critical point of  $W$  is smooth outside of a finite number of points (called branch points, where  $\vec{\Phi}$  fails to be an immersion), but globally only in  $W^{2,p}(\Sigma, \mathbb{R}^n)$  for all  $p < \infty$ . In particular, if the branch points are  $p_1, \dots, p_m \in \Sigma$ , we could have

$$(3) \quad d \left( *_g d \vec{H}_g - 3 *_g (d \vec{H}_g)^N + \star(\vec{H}_g \wedge d\vec{n}) \right) = \sum_{i=1}^m \vec{\alpha}_i \delta_{p_i}$$

for some  $\vec{\alpha}_1, \dots, \vec{\alpha}_m \in \mathbb{R}^n$ , or more generally derivatives of Dirac masses.

DEFINITION 1. – We say that a branched Willmore immersion is *variational* if it is obtained as a weak limit or as a bubble of a sequence of Willmore immersions of uniformly bounded energy.

The Equation (3) permits to introduce the first residue defined in [2] (see formula (1-8) p. 260) as

$$(4) \quad \tilde{\gamma}_0(p_i) = \frac{1}{4\pi} \int_{\gamma} *_g d \vec{H}_g - 3 *_g (d \vec{H}_g)^N + \star(\vec{H}_g \wedge d\vec{n}) = \frac{1}{2} \vec{\alpha}_i$$

for any smooth closed curved  $\gamma$  around  $p_i$ , for  $i = 1, \dots, m$ . This quantity was first defined for immersions in codimension 1 by Kuwert and Schätzle in [17] (Lemma 4.1 p. 338), and in any codimension in [2]. We will see that  $\tilde{\gamma}_0(p_i)$  measures on the basic first obstruction to the regularity of Willmore surfaces through the branch points.

DEFINITION 2. – *We say that a branched Willmore surface  $\vec{\Phi} : \Sigma \rightarrow \mathbb{R}^n$  is a true branched immersion if for all branch point  $p \in \Sigma$ , the first residue  $\tilde{\gamma}_0(p)$  vanishes (i.e.,  $\tilde{\gamma}_0(p) = 0$ ).*