

Astérisque

JAN NEKOVÁŘ

Selmer complexes

Astérisque, tome 310 (2006)

[<http://www.numdam.org/item?id=AST_2006__310__R1_0>](http://www.numdam.org/item?id=AST_2006__310__R1_0)

© Société mathématique de France, 2006, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ASTÉRIQUE 310

SELMER COMPLEXES

Jan Nekovář

Société Mathématique de France 2006

Publié avec le concours du Centre National de la Recherche Scientifique

J. Nekovář

E-mail : nekovar@math.jussieu.fr

Url : <http://www.institut.math.jussieu.fr/~nekovar/>

Université Pierre et Marie Curie (Paris 6), Institut de Mathématiques de Jussieu,
Théorie des Nombres, Case 247, 4, place Jussieu, F-75252 Paris Cedex 05, FRANCE.

2000 Mathematics Subject Classification. — 11R23, 11G40, 11F41.

Key words and phrases. — Iwasawa theory, Selmer groups, big Galois representations.

SELMER COMPLEXES

Jan Nekovář

Abstract. — This book builds new foundations of Iwasawa theory, based on a systematic study of cohomological invariants of big Galois representations in the framework of derived categories. A new duality formalism is developed, which leads to generalized Cassels-Tate pairings and generalized p -adic height pairings. One of the applications is a parity result for Selmer groups associated to Hilbert modular forms.

Résumé (Complexes de Selmer). — Ce livre construit de nouvelles fondations pour la théorie d'Iwasawa, basées sur une étude systématique d'invariants cohomologiques (vivant dans des catégories dérivées) pour les grosses représentations galoisiennes. On développe un nouveau formalisme de dualité dont on déduit des accouplements de Cassels-Tate généralisés et des hauteurs p -adiques généralisées. Une des applications est un résultat de parité pour les groupes de Selmer attachés aux formes modulaires de Hilbert.

CONTENTS

0. Introduction	1
0.0. Big Galois representations	1
0.1. Examples	1
0.2. Selmer groups	2
0.3. Big vs. finite Galois representations	3
0.4. Compact vs. discrete modules	4
0.5. (Ind-)admissible $R[G]$ -modules	6
0.6. Continuous cohomology	7
0.7. Duality for Galois cohomology	7
0.8. Selmer complexes	9
0.9. Duality for Selmer complexes	10
0.10. Comparison with classical Selmer groups	12
0.11. Iwasawa theory	13
0.12. Duality for Galois cohomology in Iwasawa theory	13
0.13. Duality for Selmer complexes in Iwasawa theory	15
0.14. Classical Iwasawa theory	16
0.15. Generalized Cassels-Tate pairings	18
0.16. Generalized height pairings	22
0.17. Parity results	25
0.18. Contents	25
0.19. Directions for further research	26
0.20. Miscellaneous	27
1. Homological Algebra: Products and Signs	29
1.1. Standard notation and conventions	29
1.2. Tensor products and Hom	33
1.3. Products	41
2. Local Duality	51
2.1. Notation	51
2.2. Dualizing functors	51
2.3. Matlis duality	52

2.4. Cohomology with support at $\{\mathfrak{m}\}$	53
2.5. Local Duality ([RD , Ch. V])	54
2.6. Grothendieck Duality ([RD , Ch. V])	55
2.7. Remarks	55
2.8. Relating D , $\mathscr{D}$ and Φ	56
2.9. Relation to Pontrjagin duality	61
2.10. Non-complete R	62
2.11. Semi-local R	74
3. Continuous Cohomology	75
3.1. Properties of R -modules of finite type	75
3.2. Admissible $R[G]$ -modules	75
3.3. Ind-admissible $R[G]$ -modules	79
3.4. Continuous cochains	82
3.5. Continuous (hyper-)cohomology	87
3.6. Derived functor cohomology	90
3.7. Localization	93
4. Continuous Cohomology of Pro-Finite Groups	95
4.1. Basic properties	95
4.2. Finiteness conditions	96
4.3. The duality diagram: T , A , T^* , A^*	101
4.4. Comparing $\mathbf{R}^+\Gamma_{\text{der}}$ and $\mathbf{R}\Gamma_{\text{cont}}$	102
4.5. Bar resolution	105
4.6. Euler-Poincaré characteristic	106
5. Duality Theorems for Galois Cohomology Revisited	113
5.1. Classical duality results for Galois cohomology	113
5.2. Duality for G_v	114
5.3. Cohomology with compact support for $G_{K,S}$	122
5.4. Duality for $G_{K,S}$	125
5.5. Duality for Poincaré groups	129
5.6. Localization	130
5.7. In the absence of (P)	130
6. Selmer Complexes	135
6.1. Definition of Selmer complexes	135
6.2. Orthogonal local conditions	137
6.3. Global cup products	140
6.4. Functoriality of Selmer complexes	143
6.5. Transpositions	145
6.6. Self-dual case	148
6.7. An example of local conditions	151
6.8. Localization	154
6.9. In the absence of (P)	154

7. Unramified cohomology	155
7.1. Notation	155
7.2. Complexes $C(M)$	156
7.3. Explicit resolutions (discrete case)	161
7.4. Explicit resolutions (pro-finite case)	163
7.5. Duality	168
7.6. From modules to complexes	171
7.7. Transpositions	181
7.8. Greenberg's local conditions	184
8. Iwasawa theory	189
8.1. Shapiro's Lemma	189
8.2. Shapiro's Lemma for ind-admissible modules	200
8.3. Infinite extensions	202
8.4. Infinite Abelian extensions	204
8.5. Duality theorems in Iwasawa theory	216
8.6. Local conditions and Shapiro's Lemma	221
8.7. Functoriality of the unramified local conditions	227
8.8. Greenberg's local conditions in Iwasawa theory	230
8.9. Duality theorems in Iwasawa theory revisited	238
8.10. Control Theorems	249
8.11. Iwasawa theory over local fields	255
8.12. In the absence of (P)	259
9. Classical Iwasawa theory	261
9.1. Generalities	261
9.2. Cohomology of $\mathbf{Z}_p(1)$	264
9.3. Pseudo-null submodules	267
9.4. Relating A'_∞ and X'_∞	270
9.5. Relating A_∞ and X_∞	281
9.6. Comparison with classical Selmer groups	286
9.7. Duality and perfectness	293
10. Generalized Cassels-Tate pairings	301
10.1. A topological analogue	301
10.2. Abstract Cassels-Tate pairings	305
10.3. Greenberg's local conditions	308
10.4. Localized Cassels-Tate pairings	320
10.5. Greenberg's local conditions - localized version	322
10.6. Discrete valuation rings with involution	324
10.7. Skew-Hermitian and symplectic pairings on generalized Selmer groups	328
10.8. Comparison with the Flach pairing	347
11. R-valued height pairings	353
11.1. Definition of the height pairing	353
11.2. Symmetry of the height pairing	357

11.3. Comparison with classical p -adic height pairings	363
11.4. Comparison with classical p -adic height pairings (bis)	375
11.5. Higher height pairings	383
11.6. Bockstein spectral sequence and descent	390
11.7. Formulas of the Birch and Swinnerton-Dyer type	398
11.8. Higher height pairings and the Cassels-Tate pairing	407
11.9. Higher height pairings and parity results	409
12. Parity of ranks of Selmer groups	413
12.1. The general setup	413
12.2. The parity results	420
12.3. Hilbert modular forms	427
12.4. Galois representations associated to Hilbert modular forms	436
12.5. Ordinary Hilbert modular forms	442
12.6. L -functions over ring class fields	455
12.7. Hida families of Hilbert modular forms	470
12.8. Level raising	498
12.9. Parity results in the dihedral case	512
12.10. Proof of Theorem 12.2.3	520
12.11. Proof of Theorem 12.2.8	528
12.12. Systematic growth of Selmer groups	537
12.13. Concluding remarks	538
<i>Errata</i>	539
List of symbols	541
Bibliography	549

CHAPTER 0

INTRODUCTION

0.0. Big Galois representations

In this work we study cohomological invariants of “big Galois representations”

$$\rho : G \longrightarrow \mathrm{Aut}_R(T),$$

where

- (i) G is a suitable Galois group.
- (ii) R is a complete local Noetherian ring, with a finite residue field of characteristic p .
- (iii) T is an R -module of finite type.
- (iv) ρ is a continuous homomorphism of pro-finite groups.

We develop a general machinery that covers duality theory, Iwasawa theory, generalized Cassels-Tate pairings and generalized height pairings.

0.1. Examples

0.1.0. An archetypal example of a big Galois representation arises as follows. Let K be a field of characteristic $\mathrm{char}(K) \neq p$. For every K -scheme $X \rightarrow \mathrm{Spec}(K)$ put $\overline{X} = X \otimes_K K^{\mathrm{sep}}$, where K^{sep} is a fixed separable closure of K . Given a projective system $X_\infty = (X_\alpha)_{\alpha \in I}$ (indexed by some directed set I) of separated K -schemes of finite type with *finite* transition morphisms $X_\beta \rightarrow X_\alpha$, put

$$H^i(X_\infty) := \varprojlim_{\alpha} H_{\mathrm{et}}^i(\overline{X}_\alpha, \mathbf{Z}_p) = \varprojlim_{\alpha} \varprojlim_n H_{\mathrm{et}}^i(\overline{X}_\alpha, \mathbf{Z}/p^n \mathbf{Z}),$$

where the transition morphisms are given by trace maps. This is a representation of $G_K = \mathrm{Gal}(K^{\mathrm{sep}}/K)$, linear over the $\mathbf{Z}_p$ -algebra generated by “endomorphisms” of the tower X_∞ . In practice, $H^i(X_\infty)$ is often too big and must be first decomposed into smaller constituents. One can also use more general coefficient sheaves, not just $\mathbf{Z}/p^n \mathbf{Z}$.