

quatrième série - tome 53 fascicule 5 septembre-octobre 2020

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Tamotsu IKEDA & Shunsuke YAMANA

On the lifting of Hilbert cusp forms to Hilbert-Siegel cusp forms

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Patrick BERNARD

Publication fondée en 1864 par Louis Pasteur

Comité de rédaction au 1^{er} janvier 2020

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE

P. BERNARD D. HARARI

de 1883 à 1888 par H. DEBRAY

S. BOUCKSOM A. NEVES

de 1889 à 1900 par C. HERMITE

G. CHENEVIER J. SZEFTTEL

de 1901 à 1917 par G. DARBOUX

Y. DE CORNULIER S. VŨ NGỌC

de 1918 à 1941 par É. PICARD

A. DUCROS A. WIENHARD

de 1942 à 1967 par P. MONTEL

G. GIACOMIN G. WILLIAMSON

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,

45, rue d'Ulm, 75230 Paris Cedex 05, France.

Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.

annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France

Case 916 - Luminy

13288 Marseille Cedex 09

Tél. : (33) 04 91 26 74 64

Fax : (33) 04 91 41 17 51

email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 428 euros.

Abonnement avec supplément papier :

Europe : 576 €. Hors Europe : 657 € (\$ 985). Vente au numéro : 77 €.

© 2020 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Fabien Durand

Périodicité : 6 n^{os} / an

ON THE LIFTING OF HILBERT CUSP FORMS TO HILBERT-SIEGEL CUSP FORMS

BY TAMOTSU IKEDA AND SHUNSUKE YAMANA

ABSTRACT. — Starting from a Hilbert cusp form of weight 2κ , we will construct a Hilbert-Siegel cusp form of weight $\kappa + \frac{m}{2}$ and degree m and its transfer to inner forms of symplectic groups. Applications include a relation between Fourier coefficients of Hilbert cusp forms of weight $n + \frac{1}{2}$ and a certain weighted sum of the representation numbers of a quadratic form of rank $2n$ by a quadratic form of rank $4n$.

RÉSUMÉ. — Partant d'une forme modulaire parabolique de Hilbert de poids 2κ , nous construisons une forme modulaire parabolique de Hilbert-Siegel de poids $\kappa + \frac{m}{2}$ et de degré m et son transfert aux formes intérieures des groupes symplectiques. Comme application, on obtient entre autres une relation entre les coefficients de Fourier de formes modulaires paraboliques de Hilbert de poids $n + \frac{1}{2}$ et une certaine somme pondérée des nombres de représentation d'une forme quadratique de rang $2n$ par une forme quadratique de rang $4n$.

1. Introduction

The present investigation deals with the following problem: starting from simple automorphic data such as cusp forms on GL_2 , construct more complicated automorphic forms on groups of higher degree. Toward this problem, Ikeda [22] has constructed a lifting associating to an elliptic cusp form a Siegel cusp form of even genus. This paper generalizes it to Hilbert cusp forms with different methods. The resulting Hilbert-Siegel cusp forms are applied to the theory of quadratic forms.

To illustrate our results, let F be a totally real number field of degree d with adèle ring $\mathbb{A}$. We write $\mathbb{A}_f$ and $\mathbb{A}_\infty$ for the finite part and the infinite part of the adèle ring. We denote the set of d real primes of F by $\mathfrak{S}_\infty$ and the normalized absolute value by $\alpha = \prod_v \alpha_v : \mathbb{A}^\times \rightarrow \mathbb{R}_+^\times$.

Let $\mathrm{Sym}_m = \{z \in \mathrm{M}_m \mid {}^t z = z\}$ be the space of symmetric matrix of size m and W_m a symplectic vector space of dimension $2m$. We take matrix representation

$$\mathrm{Sp}_m = \left\{ g \in \mathrm{GL}_{2m} \mid g \begin{pmatrix} 0 & -\mathbf{1}_m \\ \mathbf{1}_m & 0 \end{pmatrix} {}^t g = \begin{pmatrix} 0 & -\mathbf{1}_m \\ \mathbf{1}_m & 0 \end{pmatrix} \right\}$$

of the associated symplectic group $\mathrm{Sp}(W_m)$ by choosing a Witt basis of W_m . We define homomorphisms $\mathfrak{m} : \mathrm{GL}_m \rightarrow \mathrm{Sp}_m$ and $\mathfrak{n} : \mathrm{Sym}_m \rightarrow \mathrm{Sp}_m$ by

$$(1.1) \quad \mathfrak{m}(a) = \begin{pmatrix} a & 0 \\ 0 & {}^t a^{-1} \end{pmatrix}, \quad \mathfrak{n}(b) = \begin{pmatrix} \mathbf{1}_m & b \\ 0 & \mathbf{1}_m \end{pmatrix}.$$

Let $\mathrm{Mp}(W_m)_{\mathbb{A}} \rightarrow \mathrm{Sp}_m(\mathbb{A})$ be the metaplectic double cover. Denote the inverse image of $\mathrm{Sp}_m(\mathbb{A}_{\infty})$ (resp. $\mathrm{Sp}_m(\mathbb{A}_{\mathfrak{f}})$) by $\mathrm{Mp}(W_m)_{\infty}$ (resp. $\mathrm{Mp}(W_m)_{\mathfrak{f}}$).

Define the character $\mathbf{e}_{\infty} : \mathbb{C}^d \rightarrow \mathbb{C}^{\times}$ by $\mathbf{e}_{\infty}(\mathcal{Z}) = \prod_{v \in \mathfrak{S}_{\infty}} e^{2\pi\sqrt{-1}\mathcal{Z}_v}$. Let $\psi = \prod_v \psi_v$ be the additive character of $\mathbb{A}/F$ whose restriction to $\mathbb{A}_{\infty}$ is $\mathbf{e}_{\infty}|_{\mathbb{R}^d}$. We let $\mathfrak{p}$ denote a finite prime of F and do not use $\mathfrak{p}$ for an archimedean place. For $\xi \in \mathrm{Sym}_m(F)$ we define the character $\psi_{\mathfrak{f}}^{\xi} = \prod_{\mathfrak{p}} \psi_{\mathfrak{p}}^{\xi} : \mathrm{Sym}_m(\mathbb{A}_{\mathfrak{f}}) \rightarrow \mathbb{C}^{\times}$ by $\psi_{\mathfrak{f}}^{\xi}(z) = \prod_{\mathfrak{p}} \psi_{\mathfrak{p}}(\mathrm{tr}(\xi z_{\mathfrak{p}}))$. For $t \in F_v^{\times}$ there is an 8th root of unity $\gamma(\psi_v^t)$ such that for all Schwartz functions ϕ on F_v

$$\int_{F_v} \phi(x_v) \psi_v(tx_v^2) dx_v = \gamma(\psi_v^t) |2t|_v^{-1/2} \int_{F_v} \mathcal{F}\phi(x_v) \psi_v\left(-\frac{x_v^2}{4t}\right) dx_v,$$

where dx_v is the self-dual Haar measure on F_v with respect to the Fourier transform $\mathcal{F}\phi(y) = \int_{F_v} \phi(x_v) \psi_v(x_v y) dx_v$. Set $\gamma^{\psi_v}(t) = \gamma(\psi_v)/\gamma(\psi_v^t)$. We denote the set of totally positive elements of F by $F_+^{\times}$, the set of totally positive definite symmetric matrices of rank m over F by Sym_m^+ and the set of all complex symmetric matrices of size m with positive definite imaginary part by $\mathcal{H}_m$. For $t \in F^{\times}$ we write $\hat{\chi}^t = \prod_v \hat{\chi}_v^t$ for the quadratic character of $\mathbb{A}^{\times}/F^{\times}$ associated to the extension $F(\sqrt{t})/F$ and denote its restriction to the finite idèle group $\mathbb{A}_{\mathfrak{f}}^{\times}$ by $\hat{\chi}_{\mathfrak{f}}^t$. For $\ell \in \mathbb{R}^d$ we will set $|t|^{\ell} = \prod_{v \in \mathfrak{S}_{\infty}} |t|_v^{\ell_v}$.

The real metaplectic group $\mathrm{Mp}(W_m)_v$ acts on $\mathcal{H}_m$ through $\mathrm{Sp}_m(F_v)$ for $v \in \mathfrak{S}_{\infty}$. There is a unique factor of automorphy $J : \mathrm{Mp}(W_m)_v \times \mathcal{H}_m \rightarrow \mathbb{C}^{\times}$ satisfying $J(\tilde{g}_v, \mathcal{Z}_v)^2 = \det(C_v \mathcal{Z}_v + D_v)$. We here write the projection of $\tilde{g}_v$ to $\mathrm{Sp}_m(F_v)$ as $\begin{pmatrix} * & * \\ C_v & D_v \end{pmatrix}$. Let ℓ be

a tuple of d positive half integers such that $2\ell_v \equiv 2\ell_{v'} \pmod{2}$ for all $v, v' \in \mathfrak{S}_{\infty}$. We set $J_{\ell}(\tilde{g}, \mathcal{Z}) = \prod_{v \in \mathfrak{S}_{\infty}} J(\tilde{g}_v, \mathcal{Z}_v)^{2\ell_v}$ for $\tilde{g} \in \mathrm{Mp}(W_m)_{\infty}$ and $\mathcal{Z} \in \mathcal{H}_m^d$. If $\ell \in \mathbb{Z}^d$, then J_{ℓ} descends to the function on $\mathrm{Sp}_m(\mathbb{A}_{\infty}) \times \mathcal{H}_m^d$. Even when $\ell \notin \mathbb{Z}^d$, one can define it on some congruence subgroup Γ_m^{θ} of $\mathrm{Sp}_m(F)$. A Hilbert-Siegel modular form $\mathcal{F}$ of weight ℓ with respect to a congruence subgroup Γ of $\mathrm{Sp}_m(F)$ is a holomorphic function on $\mathcal{H}_m^d$ which satisfies $\mathcal{F}(\gamma \mathcal{Z}) = \mathcal{F}(\mathcal{Z}) J_{\ell}(\gamma, \mathcal{Z})$ for every $\gamma \in \Gamma$ and also the additional condition at infinity if $m = 1$ and $F = \mathbb{Q}$. Let $M_{\ell}(\Gamma)$ denote the vector space of such Hilbert-Siegel modular forms. The vector space $S_{\ell}(\Gamma)$ of Hilbert-Siegel cusp forms consists of functions $\mathcal{F} \in M_{\ell}(\Gamma)$ such that $\mathcal{F}(\gamma \mathcal{Z}) \sqrt{J_{2\ell}(\gamma, \mathcal{Z})}^{-1}$ has a Fourier expansion of the form $\sum_{\xi \in \mathrm{Sym}_m^+} A(\xi) \mathbf{e}_{\infty}(\mathrm{tr}(\xi \mathcal{Z}))$ for all $\gamma \in \mathrm{Sp}_m(F)$, where $\sqrt{J_{2\ell}(\gamma, \mathcal{Z})}$ means any branch of the square root of $J_{2\ell}(\gamma, \mathcal{Z})$. Let $S_{\ell}^{(m)}$ denote the union of $S_{\ell}(\Gamma)$ for congruence subgroups $\Gamma \subset \Gamma_m^{\theta}$. The group $\mathrm{Mp}(W_m)_{\mathfrak{f}}$ acts on the space $S_{\ell}^{(m)}$ and it is important to know which representations appear in this space. We shall explicitly construct a rather small irreducible submodule, which is neither tempered nor generic at any finite prime.

We define the space $\mathfrak{C}_{2\kappa}$ of Hilbert cusp forms on PGL_2 of weight 2κ in Definition 5.4. Let $\pi_{\mathfrak{f}} \simeq \otimes'_{\mathfrak{p}} \pi_{\mathfrak{p}}$ be an irreducible admissible unitary generic representation of $\mathrm{PGL}_2(\mathbb{A}_{\mathfrak{f}})$. For some reason (see Remark 6.3) we suppose that none of $\pi_{\mathfrak{p}}$ is supercuspidal, i.e., there is a

collection of continuous characters $\mu_{\mathfrak{p}}$ of the multiplicative groups of nonarchimedean local fields $F_{\mathfrak{p}}$ such that $\pi_{\mathfrak{f}}$ is equivalent to the unique irreducible submodule of the principal series representation $\bigotimes'_{\mathfrak{p}} I(\mu_{\mathfrak{p}}, \mu_{\mathfrak{p}}^{-1})$, where $\mu_{\mathfrak{p}}$ is unramified for almost all $\mathfrak{p}$. Put $\mu_{\mathfrak{f}} = \prod_{\mathfrak{p}} \mu_{\mathfrak{p}}$. We form the restricted tensor product $I_m^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}}) = \bigotimes'_{\mathfrak{p}} I_m^{\psi_{\mathfrak{p}}}(\mu_{\mathfrak{p}})$, where $I_m^{\psi_{\mathfrak{p}}}(\mu_{\mathfrak{p}})$ is the representation of the local metaplectic group $\mathrm{Mp}(W_m)_{\mathfrak{p}}$ on the space of smooth functions $h_{\mathfrak{p}}$ on $\mathrm{Mp}(W_m)_{\mathfrak{p}}$ transforming on the left according to

$$h_{\mathfrak{p}}((\mathbf{m}(a)\mathbf{n}(b), \zeta)\tilde{g}) = \zeta^m \gamma^{\psi_{\mathfrak{p}}}(\det a)^m \mu_{\mathfrak{p}}(\det a) |\det a|_{\mathfrak{p}}^{(m+1)/2} h_{\mathfrak{p}}(\tilde{g})$$

for all $\zeta \in \{\pm 1\}$, $a \in \mathrm{GL}_m(F_{\mathfrak{p}})$, $z \in \mathrm{Sym}_m(F_{\mathfrak{p}})$ and $\tilde{g} \in \mathrm{Mp}(W_m)_{\mathfrak{p}}$. This representation has a unique irreducible submodule $A_m^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}})$, which is unitary.

THEOREM 1.1. – *Notation being as above, $A_m^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}})$ appears in $S_{(2\kappa+m)/2}^{(m)}$ if and only if $\pi_{\mathfrak{f}}$ appears in $\mathfrak{C}_{2\kappa}$ and $(-1)^{\sum_{v \in \mathfrak{S}_{\infty}} \kappa_v} \prod_{\mathfrak{p}} \mu_{\mathfrak{p}}(-1) = 1$.*

We here denote the tuple $(\kappa_v + \frac{m}{2})_{v \in \mathfrak{S}_{\infty}} \in \frac{1}{2} \mathbb{Z}^d$ simply by $(2\kappa + m)/2$. The representation $\pi_{\mathfrak{f}}$ is the Shimura correspondence of $A_1^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}})$ and $A_2^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}})$ is the Saito-Kurokawa lifting of $\pi_{\mathfrak{f}}$. Both are theta liftings. We discuss the connection of this result with Arthur's endoscopic classification in §6.2. Though the trace formula will ultimately lead to another proof, our proof, which relies heavily on the theory of the Shimura correspondence but not on the Saito-Kurokawa lifting, is completely elementary. If $\kappa_v < m$ for every $v \in \mathfrak{S}_{\infty}$, then we obtain an irreducible cuspidal automorphic representation which is nontempered at all the places.

More importantly, our proof gives more precise information. We can describe how the representation $A_m^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}})$ is embedded in $S_{(2\kappa+m)/2}^{(m)}$ quite explicitly. Fix a Haar measure $db = \bigotimes_{\mathfrak{p}} db_{\mathfrak{p}}$ on $\mathrm{Sym}_m(\mathbb{A}_{\mathfrak{f}})$. Then we can associate to each $\xi \in \mathrm{Sym}_m^+$ a basis vector $w_{\xi}^{\mu_{\mathfrak{f}}}$ of the one-dimensional vector space $\mathrm{Hom}_{\mathrm{Sym}_m(\mathbb{A}_{\mathfrak{f}})}(I_m^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}}) \circ \mathbf{n}, \psi_{\mathfrak{f}}^{\xi})$ by

$$w_{\xi}^{\mu_{\mathfrak{f}}}(\bigotimes_{\mathfrak{p}} h_{\mathfrak{p}}) = \prod_{\mathfrak{p}} w_{\xi}^{\mu_{\mathfrak{p}}}(h_{\mathfrak{p}}),$$

where $w_{\xi}^{\mu_{\mathfrak{p}}} \in \mathrm{Hom}_{\mathrm{Sym}_m(F_{\mathfrak{p}})}(I_m^{\psi_{\mathfrak{p}}}(\mu_{\mathfrak{p}}) \circ \mathbf{n}, \psi_{\mathfrak{p}}^{\xi})$ is defined by

$$\begin{aligned} w_{\xi}^{\mu_{\mathfrak{p}}}(h_{\mathfrak{p}}) &= \int_{\mathrm{Sym}_m(F_{\mathfrak{p}})} h_{\mathfrak{p}} \left(\left(\begin{pmatrix} 0 & 1_m \\ -1_m & 0 \end{pmatrix} \mathbf{n}(b_{\mathfrak{p}}), 1 \right) \right) \overline{\psi_{\mathfrak{p}}^{\xi}(b_{\mathfrak{p}})} db_{\mathfrak{p}} \\ &\times \frac{|\det \xi|_{\mathfrak{p}}^{(m+1)/4}}{L(\frac{1}{2}, \mu_{\mathfrak{p}} \hat{\chi}_{\mathfrak{p}}^{\det \xi})} \prod_{j=1}^{[(m+1)/2]} L(2j-1, \mu_{\mathfrak{p}}^2) \times \begin{cases} 1 & \text{if } 2 \nmid m, \\ L\left(\frac{m+1}{2}, \mu_{\mathfrak{p}} \hat{\chi}_{\mathfrak{p}}^{(-1)^{m/2}}\right) & \text{if } 2 \mid m. \end{cases} \end{aligned}$$

The integral diverges but makes sense as it stabilizes. One can check that $w_{\xi}^{\mu_{\mathfrak{p}}}(h_{\mathfrak{p}}) = 1$ for almost all $\mathfrak{p}$.

THEOREM 1.2. – *If $\pi_{\mathfrak{f}}$ appears in $\mathfrak{C}_{2\kappa}$ and $(-1)^{\sum_{v \in \mathfrak{S}_{\infty}} \kappa_v} \prod_{\mathfrak{p}} \mu_{\mathfrak{p}}(-1) = 1$, then $A_m^{\psi_{\mathfrak{f}}}(\mu_{\mathfrak{f}})$ appears in the decomposition of $S_{(2\kappa+m)/2}^{(m)}$ with multiplicity one, and there is a set $\{c_t\}_{t \in F_+^{\times}}$*