

BULLETIN DE LA S. M. F.

PÁL ERDÖS

ISTVÁN JOÓ

VILMOS KOMORNIK

Characterization of the unique expansions

$1 = \sum_{i=1}^{\infty} q^{-n_i}$ and related problems

Bulletin de la S. M. F., tome 118, n° 3 (1990), p. 377-390

http://www.numdam.org/item?id=BSMF_1990__118_3_377_0

© Bulletin de la S. M. F., 1990, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (<http://smf.emath.fr/Publications/Bulletin/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

CHARACTERIZATION OF THE UNIQUE EXPANSIONS

$1 = \sum_{i=1}^{\infty} q^{-n_i}$ AND RELATED PROBLEMS

BY

PÁL ERDÖS, ISTVÁN JOÓ and VILMOS KOMORNIK (*)

RÉSUMÉ. — On caractérise les développements uniques de 1 en bases non entières. On donne une estimation pour la longueur des chiffres 0 consécutifs dans les développements gloutons. On établit certaines relations entre ces propriétés et les nombres de Pisot.

ABSTRACT. — We characterize the unique expansions in non-integer bases. We estimate the length of consecutive 0 digits in the greedy expansions. We obtain some relations between these properties and the Pisot numbers.

0. Introduction

Consider a number $1 < q < 2$. By an expansion of a real number x we mean a representation of the form

$$x = \sum_{i=1}^{\infty} \varepsilon_i q^{-i}, \quad \varepsilon_i \in \{0, 1\}.$$

It is clear that x has an expansion if and only if $0 \leq x \leq 1/(q-1)$.

Let us introduce the lexicographic order $\overset{L}{<}$ between the real sequences : $(\varepsilon_i) \overset{L}{<} (\varepsilon'_i)$ if there is a positive integer m such that $\varepsilon_i = \varepsilon'_i$ for all $i < m$ and $\varepsilon_m < \varepsilon'_m$. It is easy to verify that for every fixed $0 \leq x \leq 1/(q-1)$

(*) Texte reçu le 21 mai 1990, révisé le 21 juin 1990.

P. ERDÖS, Mathematical Institute of the Hungarian Academy of Sciences, Reáltanoda u. 13–15, 1053 Budapest, Hongrie.

I. JOÓ, Eötvös Loránd University, Dept. of Analysis, Múzeum krt. 6-8, 1088 Budapest, and Mathematical Institute of the Hungarian Academy of Sciences, Reáltanoda u. 13–15, 1053 Budapest, Hongrie.

V. KOMORNIK, Eötvös Loránd University, Dept of Analysis, Múzeum krt. 6-8, 1088 Budapest, Hongrie et Univ. de Bordeaux I, UFR de Mathématiques, 351 cours de la Libération, 33405 Talence Cedex, France.

in the set of all expansions of x there is a greatest and a smallest element with respect to this order : the so-called *greedy* and *lazy* expansion, cf. [4]. (The greedy expansions were studied earlier in [1] where they were called β -expansions.) A number x has a unique expansion if and only if its greedy and lazy expansions coincide.

Let us recall that the digits of these expansions may be defined recursively as follows : if $m \geq 1$ and if the digits ε_i of the greedy expansion of x are defined for all $i < m$, then we put

$$\varepsilon_m = \begin{cases} 1 & \text{if } \sum_{i < m} \varepsilon_i q^{-i} + q^{-m} \leq x, \\ 0 & \text{if } \sum_{i < m} \varepsilon_i q^{-i} + q^{-m} > x. \end{cases}$$

If $m \geq 1$ and if the digits ε_i of the lazy expansion of x are defined for all $i < m$, then we put

$$\varepsilon_m = \begin{cases} 0 & \text{if } \sum_{i < m} \varepsilon_i q^{-i} + \sum_{i > m} q^{-i} \geq x, \\ 1 & \text{if } \sum_{i < m} \varepsilon_i q^{-i} + \sum_{i > m} q^{-i} < x. \end{cases}$$

In section 1 we characterize the unique expansions of 1. This improves some earlier results in [5]. As a by-product we obtain a new proof for the characterization of the greedy expansions, obtained earlier in [2].

In [4] it was proved that for almost every $1 < q < 2$ the greedy expansion of 1 contains arbitrarily long sequences of consecutive 0 digits. In section 2 we improve this result by giving an explicit estimate on the length of these sequences. An analogous result is obtained for the lazy expansions, too.

In section 3 we generalize some other results obtained in [4]–[7].

At the end of this paper we formulate some open questions.

The authors wish to thank the referee for drawing their attention to the papers [2], [3] and [9].

1. Characterization of the greedy and the unique expansions of 1

Fix $1 < q < 2$ arbitrarily and consider an expansion of 1 :

$$(1) \quad 1 = \sum_{i=1}^{\infty} \varepsilon_i q^{-i}, \quad \varepsilon_i \in \{0, 1\}.$$

THEOREM 1

a) (1) is the greedy expansion of 1 if and only if

$$(2) \quad (\varepsilon_{k+i}) \stackrel{L}{<} (\varepsilon_i) \quad \text{whenever } \varepsilon_k = 0.$$

b) (1) is the unique expansion of 1 if and only if (2) and

$$(3) \quad (1 - \varepsilon_{k+i}) \stackrel{L}{<} (\varepsilon_i) \quad \text{whenever } \varepsilon_k = 1.$$

are satisfied. $\square$

Remark 1. — It is easy to deduce from this theorem that if (1) is the greedy (resp. unique) expansion of 1, then (2) (resp. (2) and (3)) is satisfied for all $k \geq 1$. $\square$

The proof of this theorem is based on some lemmas concerning the more general expansions

$$(4) \quad x = \sum_{i=1}^{\infty} \varepsilon_i q^{-i}, \quad \varepsilon_i \in \{0, 1\}$$

for arbitrarily fixed $1 < q < 2$ and $0 \leq x \leq 1/(q-1)$.

LEMMA 1

a) (4) is the greedy expansion of x if and only if

$$(5) \quad \sum_{i=1}^{\infty} \varepsilon_{k+i} q^{-i} < 1 \quad \text{whenever } \varepsilon_k = 0.$$

b) (4) is the lazy expansion of x if and only if

$$(6) \quad \sum_{i=1}^{\infty} (1 - \varepsilon_{k+i}) q^{-i} < 1 \quad \text{whenever } \varepsilon_k = 1.$$

Proof :

a) If (5) is not satisfied for some $\varepsilon_k = 0$, then x has another expansion

$$(7) \quad x = \sum_{i=1}^{\infty} \varepsilon'_i q^{-i}, \quad \varepsilon'_i \in \{0, 1\}$$

such that $\varepsilon_i = \varepsilon'_i$ for all $i < k$ and $\varepsilon'_k = 1$. Then the expansion (4) is not greedy.

If the expansion (4) is not greedy, then there is another expansion (7) of x and there is a positive integer k such that $\varepsilon_i = \varepsilon'_i$ for all $i < k$ and $\varepsilon_k = 0$, $\varepsilon'_k = 1$. It follows that

$$\sum_{i>k} \varepsilon_i q^{-i} \geq q^{-k}$$

and therefore (5) is not satisfied.

b) The assertion follows at once from a) if we remark that the expansion (4) is lazy if and only if the expansion

$$(8) \quad 1/(q-1) - x = \sum_{i=1}^{\infty} (1 - \varepsilon_i) q^{-i}$$

is greedy. $\square$

LEMME 2

a) If $x \geq 1$ and if the expansion (4) is greedy, then (2) is satisfied.

b) If $x \geq 1$ and if the expansion (4) is unique, then (2) and (3) are satisfied.

Proof :

a) Assume that (2) is not satisfied for some $\varepsilon_k = 0$, then either $(\varepsilon_{k+i}) = (\varepsilon_i)$ or $(\varepsilon_{k+i}) \overset{L}{>} (\varepsilon_i)$. In the first case we have

$$\sum_{i=1}^{\infty} \varepsilon_{k+i} q^{-i} = \sum_{i=1}^{\infty} \varepsilon_i q^{-i} = x \geq 1;$$

hence the condition (5) of LEMMA 1 is not satisfied and the expansion (4) is not greedy. In the second case there is an integer m such that $\varepsilon_{k+i} = \varepsilon_i$ for all $i < m$ and $\varepsilon_{k+m} = 1$, $\varepsilon_m = 0$. If the expansion (4) were greedy, then by LEMMA 1 we would have

$$\sum_{i=1}^{\infty} \varepsilon_{k+i} q^{-i} < 1 \leq x.$$

Therefore x would have another expansion (7) such that $\varepsilon'_i = \varepsilon_i$ for all $i < m$ and $\varepsilon'_m > \varepsilon_m$; hence $(\varepsilon'_i) \overset{L}{>} (\varepsilon_i)$. But this is impossible because (4) is the greedy expansion.