

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

ON THE IWAHORI WEYL GROUP

Timo Richarz

Tome 144
Fascicule 1

2016

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 117-124

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 1, tome 144, janvier 2016

Comité de rédaction

Valérie BERTHÉ	Marc HERZLICH
Gérard BESSON	O'Grady KIERAN
Emmanuel BREUILLARD	Julien MARCHÉ
Yann BUGAUD	Emmanuel RUSS
Jean-François DAT	Christophe SABOT
Charles FAVRE	Wilhelm SCHLAG
Raphaël KRIKORIAN (dir.)	

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 178 €, hors Europe : 194 € (\$ 291)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2016

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Marc PEIGNÉ

ON THE IWAHORI WEYL GROUP

BY TIMO RICHARZ

ABSTRACT. — Let F be a discretely valued complete field with valuation ring $\mathcal{O}_F$ and perfect residue field k of cohomological dimension ≤ 1 . In this paper, we generalize the Bruhat decomposition in Bruhat and Tits [3] from the case of simply connected F -groups to the case of arbitrary connected reductive F -groups. If k is algebraically closed, Haines and Rapoport [4] define the Iwahori-Weyl group, and use it to solve this problem. Here we define the Iwahori-Weyl group in general, and relate our definition of the Iwahori-Weyl group to that of [4].

Let F be a discretely valued complete field with valuation ring $\mathcal{O}_F$ and perfect residue field k of cohomological dimension ≤ 1 . In this paper, we generalize the Bruhat decomposition in Bruhat and Tits [3] from the case of simply connected F -groups to the case of arbitrary connected reductive F -groups. If k is algebraically closed, Haines and Rapoport [4] define the Iwahori-Weyl group, and use it to solve this problem. Here we define the Iwahori-Weyl group in general, and relate our definition of the Iwahori-Weyl group to that of [4]. Furthermore, we study the length function on the Iwahori-Weyl group, and use it to determine the number of points in a Bruhat cell, when k is a finite field. Except for Lemma 1.3 below, the results are independent of [4], and are directly based on the work of Bruhat and Tits [2], [3].

Texte reçu le 13 janvier 2014, modifié et accepté le 4 avril 2015.

TIMO RICHARZ, Timo Richarz: Mathematisches Institut der Universität Bonn, Endenicher Allee 60, 53115 Bonn, Germany • *E-mail* : richarz@math.uni-bonn.de

2010 Mathematics Subject Classification. — 02F55, 20G25.

Key words and phrases. — Affine Weyl group, Reductive groups over local fields, Bruhat decomposition.

ACKNOWLEDGEMENT. — I thank my advisor M. Rapoport for his steady encouragement and advice during the process of writing. I am grateful to the stimulating working atmosphere in Bonn and for the funding by the Max-Planck society.

Let $\bar{F}$ be the completion of a separable closure of F . Let $\check{F}$ be the completion of the maximal unramified subextension with valuation ring $\mathcal{O}_{\check{F}}$ and residue field $\bar{k}$. Let $I = \text{Gal}(\bar{F}/\check{F})$ be the inertia group of $\check{F}$, and let $\Sigma = \text{Gal}(\check{F}/F)$.

Let G be a connected reductive group over F , and denote by $\mathcal{B} = \mathcal{B}(G, F)$ the enlarged Bruhat-Tits building. Fix a maximal F -split torus A . Let $\mathcal{A} = \mathcal{A}(G, A, F)$ be the apartment of $\mathcal{B}$ corresponding to A .

1.1. Definition of the Iwahori-Weyl group. — Let $M = Z_G(A)$ be the centralizer of A , an anisotropic group, and let $N = N_G(A)$ be the normalizer of A . Denote by $W_0 = N(F)/M(F)$ the relative Weyl group.

DEFINITION 1.1. — i) The *Iwahori-Weyl group* $W = W(G, A, F)$ is the group

$$W \stackrel{\text{def}}{=} N(F)/M_1,$$

where M_1 is the unique parahoric subgroup of $M(F)$.

ii) Let $\mathfrak{a} \subset \mathcal{A}$ be a facet and $P_{\mathfrak{a}}$ the associated parahoric subgroup. The subgroup $W_{\mathfrak{a}}$ of the Iwahori-Weyl group corresponding to $\mathfrak{a}$ is the group

$$W_{\mathfrak{a}} \stackrel{\text{def}}{=} P_{\mathfrak{a}} \cap N(F)/M_1.$$

The group $N(F)$ operates on $\mathcal{A}$ by affine transformations

$$(1.1) \quad \nu : N(F) \longrightarrow \text{Aff}(\mathcal{A}).$$

The kernel $\ker(\nu)$ is the unique maximal compact subgroup of $M(F)$ and contains the compact group M_1 . Hence, the morphism (1.1) induces an action of W on $\mathcal{A}$.

Let G_1 be the subgroup of $G(F)$ generated by all parahoric subgroups, and define $N_1 = G_1 \cap N(F)$. Fix an alcove $\mathfrak{a}_C \subset \mathcal{A}$, and denote by B the associated Iwahori subgroup. Let $\mathbb{S}$ be the set of simple reflections at the walls of $\mathfrak{a}_C$. By Bruhat and Tits [3, Prop. 5.2.12], the quadruple

$$(1.2) \quad (G_1, B, N_1, \mathbb{S})$$

is a (double) Tits system with affine Weyl group $W_{\text{af}} = N_1/N_1 \cap B$, and the inclusion $G_1 \subset G(K)$ is B - N -adapted of connected type.

LEMMA 1.2. — i) *There is an equality $N_1 \cap B = M_1$.*

ii) *The inclusion $N(F) \subset G(F)$ induces a group isomorphism $N(F)/N_1 \xrightarrow{\cong} G(F)/G_1$.*

Proof. — The group $N_1 \cap B$ operates trivially on $\mathcal{A}$ and so is contained in $\ker(\nu) \subset M(F)$. In particular, $N_1 \cap B = M(F) \cap B$. But $M(F) \cap B$ is a parahoric subgroup of $M(F)$ and therefore equal to M_1 .

The group morphism $N(F)/N_1 \rightarrow G(F)/G_1$ is injective by definition. We have to show that $G(F) = N(F) \cdot G_1$. This follows from the fact that the inclusion $G_1 \subset G(F)$ is B - N -adapted, cf. [2, 4.1.2]. $\square$

Kottwitz defines in [5, §7] a surjective group morphism

$$(1.3) \quad \kappa_G : G(F) \longrightarrow X^*(Z(\hat{G})^I)^\Sigma.$$

Note that in [loc. cit.] the residue field k is assumed to be finite, but the arguments extend to the general case.

LEMMA 1.3. — *There is an equality $G_1 = \ker(\kappa_G)$ as subgroups of $G(F)$.*

Proof. — For any facet $\mathfrak{a}$, let $\text{Fix}(\mathfrak{a})$ be the subgroup of $G(F)$ which fixes $\mathfrak{a}$ pointwise. The intersection $\text{Fix}(\mathfrak{a}) \cap \ker(\kappa_G)$ is the parahoric subgroup associated to $\mathfrak{a}$, cf. [4, Proposition 3]. This implies $G_1 \subset \ker(\kappa_G)$. For any facet $\mathfrak{a}$, let $\text{Stab}(\mathfrak{a})$ be the subgroup of $G(F)$ which stabilizes $\mathfrak{a}$. Fix an alcove $\mathfrak{a}_C$. There is an equality

$$(1.4) \quad \text{Fix}(\mathfrak{a}_C) \cap G_1 = \text{Stab}(\mathfrak{a}_C) \cap G_1,$$

and (1.4) holds with G_1 replaced by $\ker(\kappa_G)$, cf. [4, Lemma 17]. Assume that the inclusion $G_1 \subset \ker(\kappa_G)$ is strict, and let $\tau \in \ker(\kappa_G) \setminus G_1$. By Lemma 1.2 ii), there exists $g \in G_1$ such that $\tau' = \tau \cdot g$ stabilizes $\mathfrak{a}_C$, and hence τ' is an element of the Iwahori subgroup $\text{Stab}(\mathfrak{a}_C) \cap \ker(\kappa_G)$. This is a contradiction, and proves the lemma. $\square$

By Lemma 1.2, there is an exact sequence

$$(1.5) \quad 1 \longrightarrow W_{\text{af}} \longrightarrow W \xrightarrow{\kappa_G} X^*(Z(\hat{G})^I)^\Sigma \longrightarrow 1.$$

The stabilizer of the alcove $\mathfrak{a}_C$ in W maps isomorphically onto $X^*(Z(\hat{G})^I)^\Sigma$ and presents W as a semidirect product

$$(1.6) \quad W = X^*(Z(\hat{G})^I)^\Sigma \ltimes W_{\text{af}}.$$

For a facet $\mathfrak{a}$ contained in the closure of $\mathfrak{a}_C$, the group $W_{\mathfrak{a}}$ is the parabolic subgroup of W_{af} generated by the reflections at the walls of $\mathfrak{a}_C$ which contain $\mathfrak{a}$.

THEOREM 1.4. — *Let $\mathfrak{a}$ (resp. $\mathfrak{a}'$) be a facet contained in the closure of $\mathfrak{a}_C$, and let $P_{\mathfrak{a}}$ (resp. $P_{\mathfrak{a}'}$) be the associated parahoric subgroup. There is a bijection*

$$\begin{aligned} W_{\mathfrak{a}} \backslash W / W_{\mathfrak{a}'} &\xrightarrow{\cong} P_{\mathfrak{a}} \backslash G(F) / P_{\mathfrak{a}'} \\ W_{\mathfrak{a}} w W_{\mathfrak{a}'} &\longmapsto P_{\mathfrak{a}} n_w P_{\mathfrak{a}'}, \end{aligned}$$

where n_w denotes a representative of w in $N(F)$.