ISSN 0012-9593

quatrième série - tome 52 *fascicule 6* *novembre-décembre 2019*

ANNALES SCIENTIFIQUES de L'ÉCOLE NORMALE SUPÉRIEURE

Françoise DAL'BO & Marc PEIGNÉ & Jean-Claude PICAUD & Andrea SAMBUSETTI

Asymptotic geometry of negatively curved manifolds of finite volume

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Patrick Bernard

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. Sainte-Claire Deville
de 1883 à 1888 par H. Debray
de 1889 à 1900 par C. Hermite
de 1901 à 1917 par G. Darboux
de 1918 à 1941 par É. Picard
de 1942 à 1967 par P. Montel

Comité de rédaction au 1er mars 2019

P. Bernard	D. Harari
S. Boucksom	A. Neves
R. Cerf	J. Szeftel
G. Chenevier	S. Vũ Ngọc
Y. de Cornulier	A. Wienhard
A. Ducros	G. Williamson

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France
Case 916 - Luminy
13288 Marseille Cedex 09
Tél. : (33) 04 91 26 74 64
Fax : (33) 04 91 41 17 51
email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 420 euros.
Abonnement avec supplément papier :
Europe : 551 €. Hors Europe : 620 € ($ 930). Vente au numéro : 77 €.

© 2019 Société Mathématique de France, Paris

En application de la loi du 1er juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).
All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Stéphane Seuret
Périodicité : 6 nos / an

Ann. Scient. Éc. Norm. Sup.
4^e série, t. 52, 2019, p. 1459 à 1485

ASYMPTOTIC GEOMETRY OF NEGATIVELY CURVED MANIFOLDS OF FINITE VOLUME

BY FRANÇOISE DAL'BO, MARC PEIGNÉ, JEAN-CLAUDE PICAUD AND ANDREA SAMBUSETTI

ABSTRACT. – We study the asymptotic behavior of simply connected Riemannian manifolds X of strictly negative curvature admitting a non-uniform lattice Γ. If the quotient manifold $\bar{X} = \Gamma\backslash X$ is asymptotically $1/4$-pinched, we prove that Γ is divergent and $U\bar{X}$ has finite Bowen-Margulis measure (which is then ergodic and totally conservative with respect to the geodesic flow); moreover, we show that, in this case, the volume growth of balls $B(x, R)$ in X is asymptotically equivalent to a purely exponential function $c(x)e^{\delta R}$, where δ is the topological entropy of the geodesic flow of $\bar{X}$. This generalizes Margulis' celebrated theorem to negatively curved spaces of finite volume. In contrast, we exhibit examples of lattices Γ in negatively curved spaces X (not asymptotically $1/4$-pinched) where, depending on the critical exponent of the parabolic subgroups and on the finiteness of the Bowen-Margulis measure, the growth function is exponential, lower-exponential or even upper-exponential.

RÉSUMÉ. – Nous décrivons le comportement asymptotique de variétés Riemanniennes simplement connexes X à courbure strictement négative, dont le groupe d'isométries contient des sous-groupes discrets Γ de co-volume fini. Plus précisément, nous montrons que lorsque la courbure est asymptotiquement 1/4-pincée, le groupe Γ est alors divergent et la mesure de Bowen-Margulis associée est finie; de plus, le volume des boules $B(x, R)$ de X est asymptotiquement équivalent à la fonction $c(x)e^{\delta R}$, où δ désigne l'exposant de Poincaré de Γ. Ce résultat généralise le célèbre théorème de Margulis au cas des réseaux non-uniformes. Nous construisons aussi toute une série d'exemples de variétés X à courbure strictement négative mais non asymptotiquement 1/4-pincée, pour lesquels le volume des boules $B(x, R)$ de X ne croît pas toujours de façon purement exponentielle.

1. Introduction

Let X be a complete, simply connected manifold with strictly negative curvature. In the sixties, G. Margulis [18], using measure theory on the foliations of the Anosov system defined by the geodesic flow, showed that if Γ is a uniform lattice of X (i.e., a torsionless, discrete

0012-9593/06/© 2019 Société Mathématique de France. Tous droits réservés
doi:10.24033/asens.2413

group of isometries such that $\bar{X} = \Gamma\backslash X$ is compact), then the *orbital function* of Γ is asymptotically equivalent [(1)] to a purely exponential function:

$$v_\Gamma(x, y, R) = \#\{\gamma \in \Gamma \mid d(x, \gamma y) < R\} \sim c_\Gamma(x, y) e^{\delta(\Gamma) R},$$

where $\delta(\Gamma) = \lim_{R\to\infty} R^{-1} \ln v_\gamma(x, x, R)$ is the *critical exponent* of Γ. By integration over fundamental domains, one then obtains an asymptotic equivalence for the *volume growth function* of X:

$$v_X(x, R) = \mathrm{vol} B(x, R) \sim m(x) e^{\delta(\Gamma) R}.$$

It is well-known that the exponent $\delta(\Gamma)$ equals the *topological entropy* of the geodesic flow of $\bar{X}$ (see [19]) and that, for uniform lattices, it is the same as the *volume entropy* $\omega(X) = \limsup \frac{1}{R} \ln v_X(x, R)$ of the manifold X. The function $m(x)$, depending on the center of the ball, is the *Margulis function* of X.

Since then, this result has been generalized in different directions. Notably, G. Knieper showed in [17] that the volume growth function of a Hadamard space X (a complete, simply connected manifold with nonpositive curvature) of rank one admitting uniform lattices is *purely exponential* [(2)], that is

$$v_X(x, R) \asymp e^{\omega(X) R}.$$

In general, he showed that $v_X(x, R) \asymp R^{\frac{d-1}{2}} e^{\omega(X) R}$ for rank d manifolds; however, as far as the authors are aware, it is still unknown whether there exists a Margulis function for Hadamard manifolds of rank 1 with uniform lattices, i.e., a function $m(x)$ such that $v_X(x, R) \sim m(x) e^{\omega(X) R}$, even in the case of surfaces. Another remarkable case is that of *asymptotically harmonic manifolds* of strictly negative curvature, where the strong asymptotic homogeneity implies the existence of a Margulis function, even without compact quotients, cp. [5].

In another direction, it seems natural to ask what happens for a Hadamard space X of negative curvature admitting *nonuniform lattices* Γ (i.e., $\mathrm{vol}(\Gamma\backslash X) < \infty$): *is v_X purely exponential and, more precisely, does X admit a Margulis function?* Let us emphasize that if X also admits a uniform lattice then X is a symmetric space of rank one (by [13], Corollary 9.2.2); therefore, we are interested in spaces which do not have uniform lattices, i.e., the universal covering of finite volume, negatively curved manifolds which are not locally symmetric.

[(1)] Given two functions $f, g : \mathbb{R}_+ \to \mathbb{R}_+$, we say that f is asymptotically equivalent to g when $\lim_{R\to+\infty} f(R)/g(R) = 1$, and we will write $f \sim g$.

[(2)] We will systematically use the following convenient notation in the paper: given two functions $f, g : \mathbb{R}_+ \to \mathbb{R}_+$, we write $f \overset{C}{\prec} g$ for $R > R_0$ (or $g \overset{C}{\succ} f$) if there exists $C > 0$ such that $f(R) \le C g(R)$ for these values of R. Similarly, $f \overset{C}{\asymp} g$ means $g \overset{C}{\prec} f \overset{C}{\prec} g$. We simply write $f \prec g$ and $f \asymp g$ when the constants C and R_0 are unessential.

The *upper* and *lower exponential growth rates* of the function f are respectively defined as:

$$\omega^+(f) = \limsup_{R\to+\infty} R^{-1} \ln f(R) \quad \text{and} \quad \omega^-(f) = \omega(f) = \liminf_{R\to+\infty} R^{-1} \ln f(R)$$

and we simply write $\omega(f)$ when the two limits coincide.

Finally, we say that f is *purely exponential* if $f \asymp e^{\omega(f) R}$, *lower-exponential* when $\limsup_{R\to+\infty} \frac{f(R)}{e^{\omega(f) R}} = 0$ and *upper-exponential* when $\liminf_{R\to+\infty} \frac{f(R)}{e^{\omega(f) R}} = +\infty$; when the weaker property $\limsup_{R\to+\infty} \frac{f(R)}{e^{\omega(f) R}} = +\infty$ holds, we say that f is *weakly upper-exponential*.

It is worth to stress here that the orbital function of Γ is closely related to the volume growth function of X, but it generally has, even for lattices, a different asymptotic behavior than $v_X(x, R)$. A precise asymptotic equivalence for v_Γ was proved by T. Roblin [22] in a very general setting, proving a dichotomy based on the finiteness of the so-called Bowen-Margulis measure associated to Γ. In order to state this dichotomy, let us recall some general definitions. The limit set of a general non elementary discrete subgroup Γ of isometries of X is the subset $\Lambda(\Gamma) \subset X(\infty)$ of accumulation points of any orbit $\Gamma \cdot x$ in X. By Patterson's construction (see [22] for a precise description), there exists on $\Lambda(\Gamma)$ a family $(\mu_x)_{x\in X}$ of finite measures, supported by $\Lambda(\Gamma)$, satisfying the following conditions: for any $x, x' \in X$ and any $g \in \Gamma$,

$$\frac{d\mu_x}{d\mu_{x'}}(\xi) = e^{-\delta(\Gamma)b_\xi(x,x')} \quad \text{and} \quad \mu_{\gamma^{-1}x} = \gamma_*\mu_x$$

where $b_\xi(x, x') = \lim_{y\to\xi} d(x, y) - d(x', y)$ is the Busemann function centered at $\xi \in \Lambda(\Gamma)$. When identifying the unit tangent bundle of X with $(X(\infty) \times X(\infty) - D) \times \mathbb{R}$ (where D denotes here the diagonal in $X(\infty) \times X(\infty)$), these two properties readily imply that the measure $e^{-\delta(\Gamma)(b_\eta(x,y)+b_\xi(x,y))}\, d\eta\, d\xi\, dt$ is a Radon measure on UX, which is invariant under the actions of both Γ and the translation flow on the third coordinate; thus, it induces a measure μ_{BM} on the unit tangent bundle $U\bar{X}$ of the quotient manifold $\bar{X} = \Gamma\backslash X$, which is invariant for the geodesic flow and called the *Bowen-Margulis measure*.

T. Roblin proved that for any non-elementary discrete group of isometries Γ of a CAT(-1) space X with non-arithmetic length spectrum [3], one has:

(a) $v_\Gamma(x, y, R) \sim c_\Gamma(x, y)e^{\delta(\Gamma)R}$ *if the measure* μ_{BM} *is finite;*

(b) $v_\Gamma(x, y, R) = o(R)e^{\delta(\Gamma)R}$, *where* $o(R)$ *is infinitesimal, otherwise.*

Thus, the behavior of $v_\Gamma(x, R)$ strongly depends on the finiteness of the measure μ_{BM}; also, the asymptotic constant can be expressed in terms of μ_{BM} and of the family of Patterson-Sullivan measures (μ_x) of Γ, as $c_\Gamma(x, y) = \frac{\|\mu_x\|\,\|\mu_y\|}{\delta(\Gamma)\cdot\|\mu_{BM}\|}$.

In this paper we restrict our attention to lattices Γ, which are fundamental examples of *geometrically finite groups*; let us describe this class. Let $C(\Gamma)$ be the convex hull in X of the limit set $\Lambda(\Gamma)$ in $X \cup X(\infty)$; the group Γ acts properly discontinuously on $C(\Gamma)$, the quotient $\bar{N}(\Gamma) = \Gamma\backslash C(\Gamma)$ is called the *Nielsen core* of $\bar{X}$. The group Γ (or the quotient manifold $\bar{X}$) is said to be *geometrically finite* when for some $\varepsilon > 0$ the ε-neighborhood $\bar{N}_\varepsilon(\Gamma)$ of $\bar{N}(\Gamma)$ has finite volume. We refer to [3] for a complete description of geometrical finiteness in variable negative curvature. Some bright examples of non-geometrically finite groups are $\mathbb{Z}^d$-covering of (compact) negatively curved manifolds, whose Bowen-Margulis measure is infinite but for which very refined counting estimates exist [21].

When Γ is a lattice, the Nielsen core $\bar{N}(\Gamma)$ equals $\bar{X}$, thus Γ is clearly geometrically finite. In section §4 we will recall a useful criterion (*Finiteness Criterion* (16), due to Dal'Bo-Otal-Peigné), to decide whether a geometrically finite group has finite Bowen-Margulis measure or not; hence, a precise asymptotics for $v_\Gamma(x, R)$ as in (a).

On the other hand, any convergent group Γ exhibits a behavior as in (b), since it certainly has infinite Bowen-Margulis measure (by Poincaré recurrence, $\mu_{BM}(U\bar{X}) < \infty$ implies that

[3] This means that the additive subgroup of $\mathbb{R}$ generated by the length of closed geodesics in $G\backslash X$ is dense in $\mathbb{R}$; it is the case, for instance, if $dim(X) = 2$, or when $G = \Gamma$ is a lattice.