ISSN 0012-9593

quatrième série - tome 53 *fascicule 2* *mars-avril 2020*

ANNALES SCIENTIFIQUES de L'ÉCOLE NORMALE SUPÉRIEURE

Robert J. BERMAN & Tamás DARVAS & Chinh H. LU

Regularity of weak minimizers of the K-energy and applications to properness and K-stability

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Patrick Bernard

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. Sainte-Claire Deville
de 1883 à 1888 par H. Debray
de 1889 à 1900 par C. Hermite
de 1901 à 1917 par G. Darboux
de 1918 à 1941 par É. Picard
de 1942 à 1967 par P. Montel

Comité de rédaction au 1er janvier 2020

P. Bernard	D. Harari
S. Boucksom	A. Neves
G. Chenevier	J. Szeftel
Y. de Cornulier	S. Vũ Ngọc
A. Ducros	A. Wienhard
G. Giacomin	G. Williamson

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France
Case 916 - Luminy
13288 Marseille Cedex 09
Tél. : (33) 04 91 26 74 64
Fax : (33) 04 91 41 17 51
email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 420 euros.
Abonnement avec supplément papier :
Europe : 551 €. Hors Europe : 620 € ($ 930). Vente au numéro : 77 €.

© 2020 Société Mathématique de France, Paris

En application de la loi du 1er juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).
All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Stéphane Seuret
Périodicité : 6 nos / an

Ann. Scient. Éc. Norm. Sup.
4 e série, t. 53, 2020, p. 267 à 289

REGULARITY OF WEAK MINIMIZERS OF THE K-ENERGY AND APPLICATIONS TO PROPERNESS AND K-STABILITY

BY ROBERT J. BERMAN, TAMÁS DARVAS AND CHINH H. LU

ABSTRACT. – Let (X, ω) be a compact Kähler manifold and $\mathscr{H}$ the space of Kähler metrics cohomologous to ω. If a csck metric exists in $\mathscr{H}$, we show that all finite energy minimizers of the extended K-energy are smooth csck metrics, partially confirming a conjecture of Y.A. Rubinstein and the second author. As an immediate application, we obtain that the existence of a csck metric in $\mathscr{H}$ implies J-properness of the K-energy, thus confirming one direction of a conjecture of Tian. Exploiting this properness result we prove that an ample line bundle (X, L) admitting a csck metric in $c_1(L)$ is K-polystable. When the automorphism group is finite, the properness result, combined with a result of Boucksom-Hisamoto-Jonsson, also implies that (X, L) is uniformly K-stable.

RÉSUMÉ. – Soient (X, ω) une variété kählérienne compacte et $\mathscr{H}$ l'espace des métriques de Kähler dans la classe de cohomologie de ω. S'il existe une métrique csck dans $\mathscr{H}$, nous montrons que tous les minimiseurs dans l'espace d'énergie finie de la fonctionnelle K-énergie sont lisses, confirmant partiellement une conjecture de Y.A. Rubinstein et le deuxième auteur. Comme une conséquence immédiate, nous en déduisons que l'existence d'une métrique csck dans $\mathscr{H}$ implique la J-propreté de la fonctionnelle K-énergie. Ceci confirme une direction de la conjecture de Tian. En utilisant cette propreté nous montrons qu'un fibré ample (X, L) admettant une métrique csck dans $c_1(L)$ est K-polystable. Quand le groupe d'automorphisme est fini le résultat de propreté, combiné avec un résultat récent de Boucksom-Hisamoto-Jonsson, implique aussi que (X, L) est uniformément K-stable.

1. Introduction and main results

Let (X, J, ω) be a compact connected Kähler manifold. By

$$\mathscr{H}_\omega = \{v \in \mathcal{C}^\infty(X) \mid \omega_v := \omega + i\partial\bar{\partial} v > 0\}$$

we denote the space of Kähler potentials. By the $\partial\bar{\partial}$-lemma of Hodge theory, up to a constant, this space is in a one-to-one correspondence with $\mathscr{H}$, the space of Kähler metrics cohomologous to ω. The problem of finding canonical metrics in $\mathscr{H}$ goes back to Calabi in the fifties. In this work we will point necessary conditions under which $\mathscr{H}$ admits constant scalar curvature Kähler (csck) metrics, in terms of energy properness.

0012-9593/02/© 2020 Société Mathématique de France. Tous droits réservés
doi:10.24033/asens.2422

We now elaborate on the terminology necessary to state our main results. To have a one-to-one correspondence between potentials and metrics, we consider the space

$$\mathscr{H}_0 := \mathscr{H}_\omega \cap \mathrm{AM}^{-1}(0),$$

and we always work on the level of potentials unless specified otherwise (for the definition of AM see (3) below). The connected Lie group of holomorphic automorphisms

$$G := \mathrm{Aut}_0(X, J)$$

acts naturally on $\mathscr{H}$ via pullbacks, hence it also acts on $\mathscr{H}_0$ (see [22, Section 5.2] for a precise description of this action on the level of potentials).

Motivated by results and ideas in conformal geometry, in the 90's Tian introduced the notion of "J-properness" on $\mathscr{H}_\omega$ [45, Definition 5.1] in terms of Aubin's nonlinear energy functional J_ω and the Mabuchi K-energy E. This condition says that for any $u_j \in \mathscr{H}_\omega$ we have

$$J_\omega(u_j) \to \infty \quad \text{implies} \quad E(u_j) \to \infty. \tag{1}$$

We refer to Section 2 for the precise definitions of J_ω and E.

Tian conjectured that existence of constant scalar curvature Kähler (csck) metrics in $\mathscr{H}_\omega$ should be equivalent to J-properness of the K-energy E [45, Remark 5.2],[47] and this was proved for Fano manifolds with G trivial [46, 50]. In [36, Theorem 1] the "strong form" of the J-properness condition (1) was obtained, confirming another conjecture of Tian from [46] (for Fano manifold with trivial G), saying that the K-energy grows at least linearly with respect to the J-functional. This stronger form has been later adopted in the literature, sometimes referred to as "coercivity".

When G is non-trivial it was known that the conjecture cannot, in general, hold as stated above and numerous modifications were proposed by Tian (see [47, Conjecture 7.12], [48]). In [22], Y.A. Rubinstein and the second named author disproved one of these conjectures, proved the remaining ones for general Fano manifolds, and the following conjecture was stated for general Kähler manifolds:

Conjecture 1.1 (Conjecture 2.8 in [22]). – *Suppose (X, ω) is a Kähler manifold. There exists a csck metric cohomologous to ω if and only if for some $C, D > 0$ we have*

$$E(u) \geq C \inf_{g \in G} J_\omega(g.u) - D, \quad u \in \mathscr{H}_\omega.$$

This "modified properness conjecture" thus reduces to Tian's original prediction in case G is trivial and was originally stated for Fano manifolds by Tian himself [48]. It was proved in this context (of Fano manifolds) in [22, Theorem 2.4], and this paper also linked the resolution of the general conjecture to a regularity question on weak minimizers of the K-energy that we elaborate now.

We denote by $(\mathscr{E}^1, d_1)$ the metric completion of $\mathscr{H}_\omega$ with respect to the L^1-type Mabuchi path length metric d_1. We refer to Sections 2.1-2.2 for more precise details about this metric structure introduced in [19]. The point of connection with the questions investigated here is the fact that d_1 metric growth is comparable to J_ω [22, Proposition 5.5], and we refer to [22, Section 4, Section 5] for a more detailed exposition on how the d_1-metric geometry relates

to J-properness. Let us now state the regularity conjecture of [22] (see [22, Conjecture 2.9]) and the theorem that connects it to Conjecture 1.1 above:

CONJECTURE 1.2 (Conjecture 2.9 in [22]). – *Suppose (X, ω) is a compact Kähler manifold. The minimizers of the extended K-energy $E : \mathcal{E}^1 \to (-\infty, +\infty]$ are smooth csck metrics.*

THEOREM 1.3 (Theorem 2.10 in [22]). – *Conjecture 1.2 implies Conjecture 1.1.*

Our first main result partially confirms Conjecture 1.2 and also a less general conjecture of X.X. Chen [17, Conjecture 6.3]:

THEOREM 1.4. – *Suppose (X, ω) is a csck manifold. If $v \in \mathcal{E}^1$ minimizes the extended K-energy $E : \mathcal{E}^1 \to (-\infty, +\infty]$, then v is a smooth csck potential. In particular there exists $g \in G$ such that $g^* \omega_v = \omega$.*

The last claim follows from the uniqueness result of [8]. Using this result and Theorem 1.3 we immediately obtain one direction of Conjecture 1.1:

THEOREM 1.5. – *Suppose (X, ω) is a csck manifold. Then for some $C, D > 0$ we have*

$$E(u) \geq C \inf_{g \in G} J_\omega(g.u) - D, \quad u \in \mathcal{H}_\omega. \tag{2}$$

The proof of Theorem 1.4 relies on the L^1-Mabuchi geometry of $\mathcal{H}_\omega$ introduced in [20, 19], the finite energy pluripotential theory of [9, 28] and the convexity methods of [22, 12] and [8]. Realizing that the metric geometry of $\mathcal{H}_\omega$ and J-properness should be related seems to have first appeared in [17, Conjecture 6.1], but this work rather proposed the use of the L^2-Mabuchi metric on $\mathcal{H}_\omega$.

As a consequence of Theorem 1.5 and the techniques of [46, 6] we obtain a result on K-polystability, originally proved by Mabuchi ([33, Main Theorem] see also [32]), using a completely different argument. Slightly less general, or different flavor results were obtained by Stoppa, Stoppa-Székelyhidi, Székelyhidi [40, Theorem 1.2], [41, Theorem 1.4], [43, Theorem A] and others. We recall the relevant terminology in the last section of the paper.

THEOREM 1.6. – *Suppose $L \to X$ is a positive line bundle. If there exists a csck metric in the class $c_1(L)$, then (X, L) is K-polystable.*

The idea of proving K-stability via properness goes back to Tian's seminal paper [46]. The main point of our approach, involving geodesic rays, is to generalize the findings of [6] from the Fano case.

In case the group G is trivial, the results in [11, 13, 24] show that properness implies uniform K-stability in the L^1-sense (for terminology, see [11, 13, 24] and references therein). Thus, as a consequence of Theorem 1.5 we obtain the following:

COROLLARY 1.7. – *Assume that (X, L) is a positive line bundle and G is trivial. If there exists a csck metric in $c_1(L)$, then (X, L) is uniformly K-stable.*