

RECONSTRUCTING A VARIETY FROM ITS TOPOLOGY
[after Kollár, building on earlier work of Lieblich and Olsson]

by Kęstutis Česnavičius

INTRODUCTION

The underlying Zariski topological space $|X|$ of an algebraic variety or, more generally, a scheme X tends to have few open subsets in comparison to topologies that underlie structures appearing in differential geometry or geometric topology. Thus, intuitively, $|X|$ is a weak invariant of X , and this intuition is confirmed by low-dimensional examples: for an algebraic curve C , the proper closed subsets of $|C|$ are the finite subsets of closed points, so $|C|$ does not see much beyond the cardinality of the algebraic closure of the base field. A more surprising example was constructed by WIEGAND and KRAUTER (1981, Cor. 1): for primes p and p' , there is a homeomorphism

$$|\mathbb{P}_{\mathbb{F}_p}^2| \simeq |\mathbb{P}_{\mathbb{F}_{p'}}^2|.$$

Topological spaces that underlie schemes (resp., affine schemes) were, in fact, completely classified by HOCHSTER (1969, Thm. 9): they are the locally spectral (resp., the spectral) topological spaces. We recall that a topological space T is *spectral* if

- it is quasi-compact and quasi-separated;
- it is *sober*: each irreducible closed $T' \subset T$ is the closure $\overline{\{t\}}$ of a unique $t \in T'$;
- the quasi-compact open subsets form a base of the topology of T .

A topological space T is *locally spectral* if it has an open cover by spectral spaces. The topological space $|X|$ of a quasi-compact and quasi-separated scheme X is spectral, so Hochster's result implies that, somewhat surprisingly, $|X|$ also underlies some affine scheme. For instance, for any field k and any $n \geq 0$, the topological space $|\mathbb{P}_k^n|$ also underlies an affine scheme (which, of course, need not be a variety over a field).

Due to the above, the recent result of KOLLÁR (2020), which is the focus of this report, came as a surprise: a projective, irreducible, normal variety X over \mathbb{C} of dimension ≥ 4 is uniquely determined by its topological space $|X|$, see Theorem 1.1 below. A resulting general expectation in this direction is captured by the following conjecture of Kollár.

Conjecture 0.1 (KOLLÁR, 2020, Conjecture 3). — For seminormal, geometrically irreducible varieties X and X' over fields k and k' , respectively, with $\text{char } k = 0$ and $\dim X \geq 2$, every homeomorphism $|X| \xrightarrow{\sim} |X'|$ underlies a scheme isomorphism $X \xrightarrow{\sim} X'$.

1. RECONSTRUCTION OF PROJECTIVE VARIETIES

The following result of Kollár builds on previous work of Lieblich and Olsson and fully resolves Conjecture 0.1 for projective, normal varieties of dimension ≥ 4 in characteristic 0. In fact, it forms the foundation of credibility for a conjecture of this sort.

Theorem 1.1 (KOLLÁR, 2020, Theorem 2). — For normal, geometrically integral, projective varieties X and X' over fields k and k' , respectively, such that either

- 1) $\dim X \geq 4$ and both k and k' are of characteristic 0; or
- 2) $\dim X \geq 3$ and both k and k' are finitely generated field extensions of \mathbb{Q} ;

every homeomorphism $|X| \xrightarrow{\sim} |X'|$ underlies a scheme isomorphism $X \xrightarrow{\sim} X'$.

Remark 1.2. Since X and X' are proper and geometrically integral, we have isomorphisms $\Gamma(X, \mathcal{O}_X) \cong k$ and $\Gamma(X', \mathcal{O}_{X'}) \cong k'$, so a scheme isomorphism $X \xrightarrow{\sim} X'$ amounts to a field isomorphism $\iota: k \xrightarrow{\sim} k'$ and an isomorphism of varieties $X \otimes_{k, \iota} k' \xrightarrow{\sim} X'$.

We will focus on case 1) because it already contains most of the main ideas while avoiding further technicalities of 2) that largely concern the Hilbert irreducibility theorem. Roughly, the proof is based on studying Weil divisors D on a normal X : such D are determined by $|X|$ alone because they may be viewed as formal \mathbb{Z} -linear combinations of the points of codimension 1 (for instance, a reduced effective divisor $D \subset X$ is the closure of a finite set of codimension 1 points in X). We will let

$$\text{Div}(X) := \bigoplus_{x \in X^{(1)}} \mathbb{Z} \quad \text{and} \quad \text{Eff}(X) := \bigoplus_{x \in X^{(1)}} \mathbb{Z}_{\geq 0}$$

denote the group of all divisors (resp., the monoid of all effective divisors) on X .

It is not clear if notions such as ampleness or linear equivalence of divisors are determined by $|X|$ alone, and the crux of the argument is in exhibiting hypotheses under which they are. The ability to topologically recognize linear equivalence eventually reduces the reconstruction problem to a combinatorial recognition theorem for projective spaces in terms of incidence of their lines and points (von Staudt's fundamental theorem of projective geometry).

A divisor D on X is *ample* if some multiple nD with $n > 0$ is a Cartier divisor whose associated line bundle $\mathcal{O}(nD)$ is ample. We let \sim denote linear equivalence of divisors and say that divisors D_1 and D_2 on X are *linearly similar*, denoted by $D_1 \sim_s D_2$, if $n_1 D_1 \sim n_2 D_2$ for some nonzero integers n_1 and n_2 . If this holds with $n_1 = n_2$, then we say that D_1 and D_2 are *\mathbb{Q} -linearly equivalent*, denoted by $D_1 \sim_{\mathbb{Q}} D_2$. When we speak of reduced (resp., irreducible) divisors, we implicitly assume that they are also effective (resp., effective and reduced). With these definitions, the overall proof of

Theorem 1.1 proceeds in the following stages, which successively reconstruct more and more of the structure of X from the topological space $|X|$, and which will be discussed individually in the indicated sections:

$$\begin{aligned} |X| &\overset{\S 2}{\rightsquigarrow} (|X|, \sim_s \text{ of irreducible ample divisors}) \\ &\overset{\S 3\text{--}\S 5}{\rightsquigarrow} (|X|, \sim \text{ of effective divisors}) \\ &\overset{\S 6}{\rightsquigarrow} X. \end{aligned}$$

The last step, namely, the determination of a normal, geometrically integral, projective variety X of dimension ≥ 2 over an infinite field from its underlying topological space $|X|$ equipped with the relation of linear equivalence between effective divisors on $|X|$ is due to LIEBLICH and OLSSON (2019).

The initial results of LIEBLICH and OLSSON (2019), although already sufficient for Theorem 1.1 above, have been sharpened and expanded in KOLLÁR, LIEBLICH, OLSSON, and SAWIN (2020).

2. RECOVERING LINEAR SIMILARITY OF AMPLE DIVISORS

NOTATION. *In this section, we let X be a normal, geometrically integral, projective variety over a field k of characteristic 0.*

The first stage of the proof of Theorem 1.1 is the reconstruction of linear similarity of irreducible ample divisors from the topological space $|X|$ alone. This requires, in particular, to be able to topologically recognize ampleness of irreducible divisors, which rests crucially on the following Lefschetz type theorem for the divisor class group.

Lemma 2.1 (RAVINDRA and SRINIVAS, 2006, Theorem 1). — *Suppose that $\dim X \geq 3$ and let \mathcal{L} be an ample line bundle on X whose linear system $\Gamma(X, \mathcal{L})$ is basepoint free. For some nonempty Zariski open $U \subset \Gamma(X, \mathcal{L})$ and every effective divisor $D \subset X$ that corresponds to a k -point in U , the following restriction map is injective:*

$$\mathrm{Cl}(X) \hookrightarrow \mathrm{Cl}(D).$$

The cited result is sharper but only applies to the base change $X_{\bar{k}}$ to an algebraic closure \bar{k} . This suffices because $\mathrm{Cl}(X) \hookrightarrow \mathrm{Cl}(X_{\bar{k}})$: to see this last injectivity, note that for any divisor H on X that represents a class in the kernel, both $\mathcal{O}(H)$ and $\mathcal{O}(-H)$ have nonzero global sections, which, since X is projective, means that $H \sim 0$.

For proving Theorem 1.1 for varieties of dimension ≤ 4 , one needs a refinement of Lemma 2.1 in which X is a surface (and D is a curve). This requires arithmetic inputs, notably a theorem of NÉRON (1952) on specialization of Picard groups. We refer to KOLLÁR (2020, Thm. 74) for this refinement of Lemma 2.1. It would also be interesting to extend Lemma 2.1 to positive characteristic because this may be useful for establishing further cases of Conjecture 0.1. For instance, we could then weaken the assumption on k in this section: we could let it be a field that is not a subfield of any $\overline{\mathbb{F}}_p$.

The following is the promised topological criterion for ampleness.

Proposition 2.2 (KOLLÁR, 2020, Lemma 67). — *Suppose that $\dim X \geq 2$. An irreducible divisor $H \subset X$ is ample if and only if for every effective divisor $D \subset X$ and distinct closed points $x, x' \in X \setminus D$, there is an effective divisor $H' \subset X$ with*

$$|H \cap D| = |H' \cap D| \quad \text{and} \quad x \in H' \text{ but } x' \notin H'.$$

Sketch of proof. — To begin with the simpler direction, we assume that H is ample, replace it by a multiple to assume that H is Cartier with associated very ample line bundle \mathcal{L} , and fix a D and $x, x' \in X \setminus D$. By EGA III₁, 2.2.4, for some $n > 0$ there is an $s \in \Gamma(X, \mathcal{L}^{\otimes n})$ that vanishes at x , does not vanish at x' , and is such that the vanishing locus of $s|_D$ is $H \cap D$. We can take H' to be the vanishing locus of s .

For the converse, we make a simplifying assumption that $\dim X \geq 3$ (for $\dim X = 2$ one needs a refinement of Lemma 2.1). To argue that H is ample, we will use Kleiman's criterion (KLEIMAN, 1966, Chap. III, Thm. 1 (i) \Leftrightarrow (iv) on p. 317), according to which it suffices to show that for all distinct closed points $x, x' \in X$, there exist an integer $n > 0$ and an effective divisor \tilde{H} such that $\tilde{H} \sim nH$ and $x \in \tilde{H}$ but $x' \notin \tilde{H}$ (this will simultaneously prove that some nH is basepoint free, so is also Cartier, as we require of ample divisors). Since X is projective, Lemma 2.1 and the Bertini theorem applied to the irreducible components of $H_{\bar{k}}$ supply a normal effective divisor $D \subset X$ not containing x, x' , or any generic point of H such that $H \cap D$ is irreducible and $\text{Cl}(X) \hookrightarrow \text{Cl}(D)$. By applying the assumption to this D , we find an effective divisor $H' \subset X$ with $|H \cap D| = |H' \cap D|$ and $x \in H'$ but $x' \notin H'$. Since $H \cap D$ is irreducible, this equality of topological spaces means that $nH|_D \sim n'H'|_D$ for some $n, n' > 0$. The injectivity of $\text{Cl}(X) \hookrightarrow \text{Cl}(D)$ then implies that $nH \sim n'H'$, and it remains to set $\tilde{H} := n'H'$. \square

Proposition 2.2 allows us to topologically recognize irreducible ample divisors on X . Granted this, the following proposition then expresses the linear similarity relation \sim_s between such divisors purely in terms of the topological space $|X|$.

Proposition 2.3 (KOLLÁR, 2020, Lemma 68). — *Suppose that $\dim X \geq 3$. Irreducible divisors $H_1, H_2 \subset X$ with H_1 ample are linearly similar if and only if for any disjoint, irreducible, closed subsets $Z_1, Z_2 \subset X$ of dimension ≥ 1 there is an irreducible divisor $H' \subset X$ with*

$$|H_1 \cap Z_1| = |H' \cap Z_1| \quad \text{and} \quad |H_2 \cap Z_2| = |H' \cap Z_2|.$$

Sketch of proof. — To begin with the simpler direction, we assume that $n_1H_1 \sim n_2H_2$ for some nonzero n_1, n_2 and fix Z_1, Z_2 as in the statement. The n_i must have the same sign: otherwise $\mathcal{O}(mH_1)$ and $\mathcal{O}(-mH_1)$ would have nonzero global sections for every large, sufficiently divisible $m > 0$. Thus, we may assume that $n_1, n_2 > 0$. After replacing n_1 and n_2 by nn_1 and nn_2 for a large $n > 0$, we then combine EGA III₁, 2.2.4 and the Bertini theorem (JOUANOLOU, 1983, 6.10) to find a global section of $\mathcal{O}(n_1H_1) \simeq \mathcal{O}(n_2H_2)$ whose vanishing locus is an irreducible ample divisor H' with the desired properties (and even such that the intersection of $H'_{\bar{k}}$ with every irreducible component of $X_{\bar{k}}$ is irreducible).

For the converse, we make a simplifying assumption that $\dim X \geq 5$ (to improve to $\dim X \geq 3$ one again needs a refinement of Lemma 2.1)—this time the assumption is more serious because the $\dim X \geq 5$ case does not suffice for Theorem 1.1. Letting H_1, H_2 be irreducible ample divisors as in the statement, we iteratively apply Lemma 2.1 (with the Bertini theorem) to build disjoint, irreducible, normal closed subschemes $Z_1, Z_2 \subset X$ that are complete intersections of dimension 2 such that $H_1 \cap Z_1 \subset Z_1$ and $H_2 \cap Z_2 \subset Z_2$ are irreducible divisors and the following restriction maps are injective:

$$\mathrm{Cl}(X) \hookrightarrow \mathrm{Cl}(Z_1) \quad \text{and} \quad \mathrm{Cl}(X) \hookrightarrow \mathrm{Cl}(Z_2).$$

Since the intersections $H_1 \cap Z_1$ and $H_2 \cap Z_2$ are irreducible, these injections and the displayed equalities involving H' ensure that $n_1 H_1 \sim n'_1 H'$ and $n_2 H_2 \sim n'_2 H'$ for some $n_i, n'_i > 0$. It then follows that $n_1 n'_2 H_1 \sim n'_1 n_2 H_2$, so that H_1 and H_2 are linearly similar, as desired. \square

Propositions 2.2 and 2.3 jointly carry out the first reconstruction step promised in §1:

$$|X| \rightsquigarrow (|X|, \sim_s \text{ of irreducible ample divisors}).$$

They also topologically determine complete intersection subvarieties as follows.

Corollary 2.4. — *Suppose that $\dim X \geq 3$ and let $H \subset X$ be an irreducible ample divisor. The topological space $|X|$ alone determines the collection of those closed subsets $Z \subset |X|$ that are set-theoretic complete H -intersections, i.e., for which there are irreducible divisors $H_i \sim_s H$ for $i = 1, \dots, r$ with $r = \mathrm{codim}(Z, X)$ such that*

$$Z = |H_1 \cap \dots \cap H_r|.$$

Proof. — Propositions 2.2 and 2.3 imply that $|X|$ alone determines the property of H being ample, as well as the linear similarity relation $H_i \sim_s H$. \square

We will call such a closed subscheme $H_1 \cap \dots \cap H_r \subset X$ a *complete H -intersection*. The requirement that the H_i be irreducible and only linearly similar (as opposed to linearly equivalent) to H makes this definition slightly nonstandard, but it is convenient because Propositions 2.2 and 2.3 only concern irreducible divisors. Any positive-dimensional complete H -intersection $H_1 \cap \dots \cap H_r$ is automatically geometrically connected by the Lefschetz hyperplane theorem (SGA 2_{new}, XII, 3.5), and the same then also holds for set-theoretic complete H -intersections.

3. RECOVERING \mathbb{Q} -LINEAR EQUIVALENCE OF AMPLE DIVISORS

NOTATION. *In this section, we let X be a normal, geometrically integral, projective variety over field k of characteristic 0 and let $H \subset X$ be an irreducible ample divisor.*

To prepare for topological recognition of linear equivalence of divisors, for now we continue to restrict to irreducible ample divisors and show how to recognize \mathbb{Q} -linear equivalence between them. This refines the result presented in the previous section because \mathbb{Q} -linear equivalence $\sim_{\mathbb{Q}}$ is a finer relation than linear similarity \sim_s . In addition, it involves techniques that will also be relevant later, such as topological

recognition of reduced 0-dimensional intersections that we discuss in Proposition 3.5 below. A key notion behind these techniques is that of (topological) linkage defined as follows.

Definition 3.1. Let $Y_1, Y_2 \subset X$ be integral closed subschemes with $\dim(Y_1 \cap Y_2) = 0$. Irreducible divisors $H_1, H_2 \subset X$ with $H_1 \sim_s H_2 \sim_s H$ are *H-linked for Y_1 and Y_2* if some irreducible divisor $\tilde{H} \subset X$ with $\tilde{H} \sim_s H$ satisfies

$$|\tilde{H} \cap Y_1| = |H_1 \cap Y_1| \quad \text{and} \quad |\tilde{H} \cap Y_2| = |H_2 \cap Y_2|.$$

The *H-linking of Y_1 and Y_2 is free* if, for some finite set of closed points $\Sigma \subset Y_1 \cup Y_2$, all H_1 and H_2 as above that are disjoint from Σ are *H-linked for Y_1 and Y_2* .

By Propositions 2.2 and 2.3, if $\dim X \geq 3$, then these notions depend only on $|X|$. They topologically encode reducedness of 0-dimensional schematic intersections as follows.

Proposition 3.2 (KOLLÁR, 2020, Proposition 81). — *Let Y_1 and Y_2 be as in Definition 3.1 and suppose that $\dim X \geq 3$, $\dim Y_1 \geq 2$, $\dim Y_2 \geq 1$, and Y_1 is geometrically connected (for instance, a set-theoretic complete *H*-intersection). Then the *H-linking of Y_1 and Y_2 is free if and only if $Y_1 \cap Y_2$ is reduced with $\Gamma(Y_i, \mathcal{O}) \xrightarrow{\sim} \Gamma(Y_1 \cap Y_2, \mathcal{O})$ for some i .**

Sketch of proof. — Since $\mathcal{O}(H)$ is ample, its global sections on $Y_1 \cup Y_2$ lift to X after possibly replacing them by powers. The H_i and \tilde{H} that appear in the definition of free *H-linking* correspond to some $s_i \in \Gamma(X, \mathcal{O}(n_i H))$ and $\tilde{s} \in \Gamma(X, \mathcal{O}(\tilde{n} H))$. Thus, in essence, the question of free *H-linking* of Y_1 and Y_2 is that of patching the sections $s_i|_{Y_i}$ along $Y_1 \cap Y_2$ to glue some of their powers to a section over $Y_1 \cup Y_2$. We may adjust the $s_i|_{Y_i}$ by global units, so the glueing is intimately related to the restriction map

$$\Gamma(Y_1, \mathcal{O})^\times \times \Gamma(Y_2, \mathcal{O})^\times \longrightarrow \Gamma(Y_1 \cap Y_2, \mathcal{O})^\times.$$

The analysis of this map eventually gives the claim, see *loc. cit.* for details. \square

Remark 3.3. The geometric connectedness of Y_1 ensures that $k \xrightarrow{\sim} \Gamma(Y_1, \mathcal{O})$ in this proposition. For X of dimension ≥ 4 , this then leads to a topological criterion for recognizing when a k -smooth closed point $x \in X$ is k -rational because one may realize such an x as the schematic intersection $Y_1 \cap Y_2$ of set-theoretic complete *H*-intersections Y_i with $\dim Y_1 = 2$ and $\dim Y_2 = 1$, see KOLLÁR (2020, Cor. 82) and Corollary 2.4 above.

More generally, by building on the idea of analyzing free *H-linking* of set-theoretic complete *H*-intersections, Kollár is able to topologically recognize isomorphy of 0-dimensional closed subschemes lying in the smooth locus as follows.

Proposition 3.4 (KOLLÁR, 2020, §84). — *Suppose that $\dim X \geq 4$ and let Z_1 and Z_2 be 0-dimensional, reduced closed subschemes of X^{sm} . Then $(|X|, \sim_s$ of irreducible ample divisors) alone determines whether or not Z_1 and Z_2 are isomorphic as k -schemes.*

In the setting of Proposition 3.2, it is also possible to topologically determine when $Y_1 \cap Y_2$ is reduced (without the additional condition on the global sections) as follows.

Proposition 3.5. — Suppose that $\dim X \geq 3$ and let $Y \subset X$ be an irreducible, geometrically connected closed k -subvariety. For irreducible, geometrically connected closed k -subvarieties $Y', Z \subset X$ such that

$$|Y \cap Z| = |Y' \cap Z| \quad \text{and this intersection is 0-dimensional,}$$

consider the following condition (that is topological by Propositions 2.2 and 2.3):

(★) $\left\{ \begin{array}{l} \text{There is a finite set of closed points } \Sigma \subset Y \cup Y' \cup Z \text{ such that all irreducible} \\ \text{divisors } H_1, H_2 \subset X \text{ disjoint from } \Sigma \text{ with } H_i \sim_s H \text{ that are } H\text{-linked for } Y' \\ \text{and } Z \text{ are also } H\text{-linked for } Y \text{ and } Z. \end{array} \right.$

1) (KOLLÁR, 2020, Cor. 89). If $\dim Y \geq 2$ and Z satisfies $\dim X - 3 \geq \dim Z \geq 1$ and is such that (★) holds for every 2-dimensional, irreducible complete H -intersection Y' , then $Y \cap Z$ is reduced.

2) (KOLLÁR, 2020, Cor. 93). If $\dim Y \geq 3$ and $y \in Y$ is a closed point such that X is k -smooth at y , then Y is k -smooth at y if and only if there is an irreducible complete H -intersection $Z \subset X$ of codimension $\dim Y$ such that $Y \cap Z$ is 0-dimensional, contains y , and (★) holds whenever Y' is an irreducible complete H -intersection.

In 1), if $Y \cap Z$ is reduced, then $Y \cap Z \subset Y' \cap Z$, so a patching of global sections of powers of $\mathcal{O}(H)$ that gives rise to an H -linking of H_1 and H_2 for Y' and Z also gives a required patching with Y in place of Y' (compare with the sketch of proof for Proposition 3.2). Thus, the main part is the converse, for which we refer to *loc. cit.* The role of the assumption on $\dim Z$ is to ensure, via the Bertini theorem, that there are many possible Y' with $|Y' \cap Z| = |Y \cap Z|$: the intersection of all such Y' is $(Y \cap Z)^{\text{red}}$.

In addition to topologically recognizing k -points and reducedness of 0-dimensional intersections as above, Kollár determines equality of intersection numbers as follows.

Proposition 3.6 (KOLLÁR, 2020, Corollary 96). — Suppose that $\dim X \geq 2$. For prime divisors $D_1, \dots, D_n \subset X$ and rational numbers $q_{ij} \in \mathbb{Q}_{>0}$ with $1 \leq i, j \leq n$, we have

$$D_i \cdot H^{\dim X - 1} = q_{ij} \cdot D_j \cdot H^{\dim X - 1} \quad \text{for all } 1 \leq i, j \leq n$$

if and only if for some closed $Z_0 \subset X$ of codimension ≥ 2 and every closed $Z \subset X$ of codimension ≥ 2 containing Z_0 there is a 1-dimensional, irreducible complete H -intersection $C \subset X$ disjoint from Z such that each $C \cap D_i$ is a disjoint union of m_i copies of $\text{Spec}(K)$ for a finite field extension K/k that does not depend on i , and $q_{ij} = m_i/m_j$.

Sketch of proof. — The ‘only if’ follows from the definitions: indeed, the intersection number $D_i \cdot H^{\dim X - 1}$ is read off from the schematic intersection $C \cap D_i$. For the ‘if,’ one first reduces to X being a surface by cutting it by irreducible ample divisors that are linearly similar to H and constructed via the Bertini theorem. Then the D_i are curves and one seeks a C cut out by some nonzero $s \in H^0(X, \mathcal{O}(mH))$ for a large $m > 0$ (such an s would lift to a section of $\mathcal{O}(mH)$ defined over the original X).

By considering $\mathcal{O}(mH)|_{\cup_{i=1}^n D_i}$ instead, it suffices to find a global section s_0 of this sheaf for a large $m > 0$ such that $\{s_0 = 0\}$ is disjoint from Z and $D_i \cap D_j$ for distinct

D_i and D_j and is a union of copies of $\text{Spec}(K)$ for some finite extension K/k . The key input to finding this s_0 is a variant of a result of POONEN (2001), according to which, for any quasi-finite, generically étale morphism $\pi: \bigcup_{i=1}^n D_i \rightarrow \mathbb{P}_k^1$, there are infinitely many separable closed points $p \in \mathbb{P}_k^1$ whose π -preimage is a reduced disjoint union of copies of p . \square

In practice, Z_0 is the nonsmooth locus $X \setminus X^{\text{sm}}$. However, $X \setminus X^{\text{sm}}$ need not *a priori* be determined by the topological space $|X|$, so, to get around this, one allows larger Z while still retaining the smoothness of $X \setminus Z$. In particular, Propositions 3.4 and 3.5.1 show that, for X of dimension ≥ 4 , Proposition 3.6 gives a purely topological criterion for determining the ratios between the intersection numbers $D_i \cdot H^{\dim X - 1}$.

Corollary 3.7. — *Suppose that $\dim X \geq 4$. For prime divisors $\{D_\lambda\}_{\lambda \in \Lambda}$ on X , the topological space $|X|$ alone determines the ratios between the $D_\lambda \cdot H^{\dim X - 1}$.*

Sketch of proof. — It suffices to apply the reasoning above to every pair $\{D_\lambda, D_{\lambda'}\}$. \square

In a similar vein, Proposition 3.6 implies the following criterion for recognizing \mathbb{Q} -linear equivalence of irreducible ample divisors.

Corollary 3.8 (KOLLÁR, 2020, Claim 97.5). — *Suppose that $\dim X \geq 2$. For irreducible ample divisors $H_1, H_2 \subset X$, we have*

$$H_1 \sim_{\mathbb{Q}} H_2 \text{ if and only if both } H_1 \sim_s H_2 \text{ and } H_1 \cdot H^{\dim X - 1} = H_2 \cdot H^{\dim X - 1}.$$

Sketch of proof. — The ‘only if’ is clear because intersection numbers are insensitive to linear equivalence. For the ‘if,’ suppose that $n_1 H_1 \sim n_2 H_2$ and use Proposition 3.6 to find a 1-dimensional, irreducible complete H -intersection $C \subset X$ such that $C \cap H_1$ and $C \cap H_2$ are 0-dimensional, reduced, and k -isomorphic. This k -isomorphism and the assumed equality of intersection numbers imply that $n_1 = n_2$, so that $H_1 \sim_{\mathbb{Q}} H_2$. \square

In the case when $\dim X \geq 4$, the criterion given by Corollary 3.8 is topological thanks to Corollary 3.7 and Propositions 2.2 and 2.3.

4. TOPOLOGICAL PENCILS OF DIVISORS

NOTATION. *In this section, we let X be a geometrically normal, geometrically integral, positive-dimensional projective variety over a field k .*

The basic idea for topologically recognizing linear equivalence between general divisors on X is to first make them ample by adding a multiple of some ample divisor and to then place them into linear pencils with a common member whose general members are irreducible. In some sense this strategy achieves a reduction to the case of irreducible ample divisors considered in §§2–3, and the key for carrying it out is to topologically describe families of divisors that end up constituting the desired pencils. The central notion is that of a topological pencil that we are going to examine in this section.

Definition 4.1. For a topological space T , a *base locus* of an infinite collection $\mathcal{P} = \{D_\lambda\}_{\lambda \in \Lambda}$ of subsets $D_\lambda \subset T$, denoted by $\text{Base}(\mathcal{P})$, is a closed subset $B \subset T$ such that, for some $\Lambda' \subset \Lambda$ whose complement is finite, we have

$$D_\lambda \cap D_{\lambda'} \subseteq B \quad \text{for all distinct } \lambda, \lambda' \in \Lambda, \text{ with } D_\lambda \cap D_{\lambda'} = B \text{ when } \lambda, \lambda' \in \Lambda'.$$

Due to the last requirement, the base locus $\text{Base}(\mathcal{P})$ is unique if it exists.

Definition 4.2 (KOLLÁR, 2020, Def. 105). A *topological pencil* on X is a set $\mathcal{P} = \{D_\lambda\}_{\lambda \in \Lambda}$ of reduced divisors $D_\lambda \subset X$ such that

- all but finitely many D_λ are irreducible;
- the D_λ jointly cover all the closed points of X ;
- $B := \text{Base}(\mathcal{P})$ exists, is of codimension ≥ 2 in X , and each $D_\lambda \setminus B$ is connected.

A topological pencil $\{D_\lambda\}_{\lambda \in \Lambda}$ is *ample* if all but finitely many D_λ are ample divisors.

Since B is of codimension ≥ 2 , the divisors D_λ in a topological pencil are connected and pairwise have no common irreducible components. The notions of a topological pencil and of its base depend only on the topological space $|X|$ and, if $\dim X \geq 2$, then, by Proposition 2.2, so does the ampleness of such a pencil. The following is the principal source of topological pencils.

Example 4.3. Let C be a normal, projective, integral k -curve and let $\pi: X \dashrightarrow C$ be a dominant rational k -morphism whose maximal locus of definition is $X \setminus B$ for a closed subset $B \subset X$. Since X is normal and C is projective and nonsingular, B is of codimension ≥ 2 in X and $\pi|_{X \setminus B}$ is flat. Let $\mathcal{P} = \{D_\lambda\}$ be the collection of the closures in X of the reduced connected components of $(\pi|_{X \setminus B})^{-1}(c)$ for a variable closed point $c \in C$. We will now show that \mathcal{P} is a topological pencil on X with base locus B . The rational map π factors through the normalization \tilde{C} of C in the maximal algebraic subextension of $k(X)/k(C)$ and this factorization has the same maximal locus of definition $X \setminus B$. By replacing C by \tilde{C} , we do not change \mathcal{P} and may assume that $k(C)$ is algebraically closed in $k(X)$, so that, by EGA IV₂, 4.5.9, the generic fiber of π is geometrically irreducible. It then follows from EGA IV₃, 9.7.7 that all but finitely many D_λ are irreducible.

Consider the closure $\bar{X} \subset X \times_k C$ of the graph of $\pi|_{X \setminus B}$, which inherits k -morphisms

$$\begin{array}{ccc} & \bar{X} & \\ b \swarrow & & \searrow \bar{\pi} \\ X & \overset{\pi}{\dashrightarrow} & C \end{array}$$

such that b is an isomorphism over $X \setminus B$ and both $\bar{\pi}$ and b are proper. The locus U of X over which b has finite fibers is open (SP, 01TI), so, since X is normal and b is birational, the finite map $b|_{b^{-1}(U)}$ is an isomorphism. In particular, $U = X \setminus B$ and B

consists precisely of the points $x \in X$ such that $\dim(b^{-1}(x)) > 0$. Since $\bar{X} \subset X \times_k C$, for closed such x the map $b^{-1}(x) \rightarrow C$ is finite surjective and $\dim(b^{-1}(x)) = 1$.

The closure in X of a reduced connected component of $(\pi|_{X \setminus B})^{-1}(c)$ is the b -image of the closure in \bar{X} of the corresponding reduced connected component of $(\bar{\pi}|_{\bar{X} \setminus b^{-1}(B)})^{-1}(c)$. Moreover, since C is nonsingular, $\bar{\pi}$ is flat and its closed fibers are purely of dimension $\dim X - 1$. Thus, we conclude from the previous paragraph that the D_λ jointly cover all the closed points of X and that the base locus of \mathcal{P} is precisely B (in Definition 4.1 choose Λ' to consist of those closed points $c \in C$ such that the c -fibers of π and $\bar{\pi}$ are geometrically irreducible).

Definition 4.4. A topological pencil \mathcal{P} on X is *algebraic* (resp., *rational*; resp., *linear*) if it is associated to some C and π as in Example 4.3 (resp., with $C_{\bar{k}} \simeq \mathbb{P}_{\bar{k}}^1$; resp., with $C \simeq \mathbb{P}_k^1$); an algebraic \mathcal{P} is *noncomposite* if $\tilde{C} = C$ in Example 4.3, that is, if every finite morphism $C' \rightarrow C$ of normal, projective, integral k -curves through which π factors is an isomorphism, equivalently, if $k(C)$ is algebraically closed in $k(X)$.

The following example relates topological pencils and linear equivalence.

Example 4.5. Let \mathcal{L} be a line bundle on X and let $s_0, s_\infty \in H^0(X, \mathcal{L})$ be nonzero global sections. The vanishing loci $D_0 := \{s_0 = 0\}$ and $D_\infty := \{s_\infty = 0\}$ are linearly equivalent divisors on X . If D_0 and D_∞ have no common irreducible components, then s_0 and s_∞ span a linear pencil whose base locus is $\{s_0 = s_\infty = 0\}$: indeed, to relate to Example 4.3, it suffices to note that s_0 and s_∞ determine a rational map $\pi: X \dashrightarrow \mathbb{P}_k^1$ whose maximal locus of definition is $X \setminus \{s_0 = s_\infty = 0\}$.

Conversely, for any pair of k -points $c, c' \in \mathbb{P}_k^1$, there is a rational function $f \in k(t)$ that vanishes to order one at c and has a simple pole at c' , so any two k -fibers of a linear topological pencil \mathcal{P} are linearly equivalent on X .

To utilize this example, we need to recognize linearity of algebraic pencils.

Lemma 4.6 (KOLLÁR, 2020, Lem. 109). — *Suppose that k is perfect and let \mathcal{P} be an algebraic topological pencil on X .*

- 1) *If $\text{Base}(\mathcal{P})$ meets the k -smooth locus $X^{\text{sm}} \subset X$, then \mathcal{P} is rational.*
- 2) *If \mathcal{P} is rational and $X^{\text{sm}}(k) \neq \emptyset$, then \mathcal{P} is linear.*
- 3) *If \mathcal{P} is rational and the k -smooth locus of some $D \in \mathcal{P}$ contains a nonempty open of some geometrically irreducible closed k -subvariety $Y \subset \text{Base}(\mathcal{P})$, then \mathcal{P} is linear.*

Proof. — Let C, π , and \bar{X} be as in Example 4.3 and let $\tilde{\bar{X}}$ be the normalization of \bar{X} . The role of the perfectness of k is to ensure that the normal k -curve C is k -smooth.

1) By a result of Abhyankar (KOLLÁR, 1996, VI.1.2), the positive dimensional fibers of a proper modification $Y' \rightarrow Y$ of excellent, normal schemes with Y regular contain nonconstant rational curves (BHATT, 2012, Rem. 4.5). We apply this to the restriction of the morphism $\tilde{\bar{X}} \rightarrow X$ to X^{sm} : by using the assumption on $\text{Base}(\mathcal{P})$, we conclude that each positive-dimensional fiber of $\tilde{\bar{X}} \rightarrow X$ receives a nonconstant morphism

from a rational curve. It then follows from Example 4.3 that $C_{\bar{k}}$ also receives such a morphism, so that $C_{\bar{k}} \simeq \mathbb{P}_{\bar{k}}^1$. Thus, \mathcal{P} is rational, as desired.

2) If $X^{\text{sm}}(k) \neq \emptyset$, then the Lang–Nishimura theorem (POONEN, 2017, 3.6.11) implies that $C(k) \neq \emptyset$. Since, by assumption, $C_{\bar{k}} \simeq \mathbb{P}_{\bar{k}}^1$, we then conclude that $C \simeq \mathbb{P}_k^1$.

3) Suppose that D arises from a closed point $c \in C$ as in Example 4.3. It suffices to argue that $[k(c) : k] = 1$, since then $C(k) \neq \emptyset$ and $C \simeq \mathbb{P}_k^1$ (compare with 2)). Since k is perfect, Example 4.3 applied over \bar{k} shows that $D_{\bar{k}}$ is a $\text{Gal}(\bar{k}/k)$ -orbit of $[k(c) : k]$ closures of connected components of fibers of π over \bar{k} . Thus, if D is k -smooth at the generic point of some geometrically irreducible k -subvariety $Y \subset \text{Base}(\mathcal{P})$, then the closures of distinct connected components of fibers of π over \bar{k} cannot simultaneously contain a nonempty open of $Y_{\bar{k}}$, so $[k(c) : k] = 1$. \square

We turn to the key question of topologically recognizing when a topological pencil is algebraic. The most basic example is the following case of an empty base locus.

Example 4.7. Suppose that k is algebraically closed. Then, by BOGOMOLOV, PIRUTKA, and SILBERSTEIN (2016, Thm. 1.1), every topological pencil $\{D_\lambda\}_{\lambda \in \Lambda}$ whose base locus is empty, in other words, such that $D_\lambda \cap D_{\lambda'} = \emptyset$ for $\lambda \neq \lambda'$, is algebraic. Indeed, *loc. cit.* says that there are a smooth, projective k -curve C and a surjective k -morphism $\pi : X \rightarrow C$ with connected fibers such that each D_λ is contained in a (closed) fiber of π . Since the D_λ jointly cover the closed points of X , the set of closed fibers of π is then precisely $\{D_\lambda\}_{\lambda \in \Lambda}$.

To proceed beyond empty base loci, it is useful to first note that algebraic topological pencils are determined by infinitely many members as follows. In essence, this is the basic reduction mechanism for reaching irreducible ample divisors from general divisors.

Lemma 4.8. — For topological pencils \mathcal{P} and \mathcal{P}' on X with \mathcal{P}' algebraic, if the set $\mathcal{P} \cap \mathcal{P}'$ of those divisors $D \subset X$ that belong to both \mathcal{P} and \mathcal{P}' is infinite, then $\mathcal{P} = \mathcal{P}'$.

Proof. — The infinitude of $\mathcal{P} \cap \mathcal{P}'$ implies that $\text{Base}(\mathcal{P}) = \text{Base}(\mathcal{P}')$, so we let B be this common base locus and let $\pi : X \dashrightarrow C$ be a dominant rational morphism that gives rise to \mathcal{P}' as in Example 4.3. As in that example, $X \setminus B$ is the maximal locus of definition of π and we may assume that the generic fiber of π is geometrically irreducible. The complements $D \setminus B$ for $D \in \mathcal{P}$ are connected and pairwise disjoint, so the infinitude of $\mathcal{P} \cap \mathcal{P}'$ ensures that each $D \setminus B$ lies in a single fiber of π . Since topological pencils cover the closed points of X , it then follows that $D \in \mathcal{P}'$ and that $\mathcal{P} = \mathcal{P}'$, as desired. \square

The following is a topological criterion for algebraicity of topological pencils.

Proposition 4.9 (KOLLÁR, 2020, Proposition 107). — Suppose that k is infinite. A topological pencil $\mathcal{P} = \{D_\lambda\}_{\lambda \in \Lambda}$ on X is algebraic if and only if for some infinite subset $\Lambda' \subset \Lambda$ and every (or merely some) irreducible ample divisor $H \subset X$, the intersection numbers

$$D_\lambda \cdot H^{\dim X - 1} \quad \text{are all equal for } \lambda \in \Lambda';$$

thus, if $\text{char}(k) = 0$ and $\dim(X) \geq 4$, then the algebraicity of \mathcal{P} depends only on $|X|$.

Sketch of proof. — The last assertion that concerns topological recognition of algebraicity of \mathcal{P} follows from the rest and from Proposition 2.2 and Corollary 3.7.

For the rest, when $\dim(X) = 1$, every \mathcal{P} is algebraic and the claim is that X has infinitely many closed points of the same degree over k . This holds because there is a finite k -morphism $X \rightarrow \mathbb{P}_k^1$ and $\mathbb{P}^1(k)$ is infinite. Thus, we may assume that $\dim X \geq 2$.

We begin with the simpler ‘only if’ direction and assume that \mathcal{P} is algebraic, associated to a $\pi: X \dashrightarrow C$ as in Example 4.3 such that all but finitely many fibers of π are geometrically irreducible. By the 1-dimensional case, there is an infinite set Λ' of closed points of C of the same degree over k with irreducible π -fibers. By using the Bertini theorem, for every irreducible ample divisor $H \subset X$ and every $\lambda, \lambda' \in \Lambda'$ we may find a complete H -intersection curve $Y \subset X$ for which $Y \cap \text{Base}(\mathcal{P}) = \emptyset$, both $Y \cap D_\lambda$ and $Y \cap D_{\lambda'}$ are 0-dimensional, and Y is flat over a neighborhood of λ and λ' in C . Then

$$D_\lambda \cdot H^{\dim X - 1} = D_{\lambda'} \cdot H^{\dim X - 1}$$

because both these intersection numbers are equal to the product of the degree of Y over C with the common degree of the points in Λ' .

For the converse, we fix a single H and let d be the common value $D_\lambda \cdot H^{\dim X - 1}$ for $\lambda \in \Lambda'$ and consider the Chow k -scheme $\text{Chow}_{X,d}$ that parametrizes those effective divisors $D \subset X$ that satisfy $D \cdot H^{\dim X - 1} = d$, so that $\text{Chow}_{X,d}$ is projective over k (KOLLÁR, 1996, Chap. I). The D_λ for $\lambda \in \Lambda'$ give infinitely many closed points on $\text{Chow}_{X,d}$, so, since $\text{Chow}_{X,d}$ is of finite type over k , their closure contains a positive-dimensional irreducible closed subscheme $C \subset \text{Chow}_{X,d}$. Consider the universal family of divisors $\pi: E \rightarrow C$ base changed from $\text{Chow}_{X,d}$, as well as the resulting commutative diagram

$$\begin{array}{ccc} E & \xrightarrow{i} & X \\ \downarrow \pi & & \\ \text{Chow}_{X,d} \supset C & & \end{array}$$

for which the D_λ for $\lambda \in \Lambda'$ appear as fibers of π . It suffices to argue that there is a nonempty open $X^0 \subset X$ such that the $i^{-1}(x)$ for a dense set of closed points $x \in X^0$ are singletons. Then, up to a power of Frobenius if $\text{char } k > 0$, the map i will be birational, C will be a curve by counting dimensions, π will give rise to an algebraic topological pencil \mathcal{P}' on X as in Example 4.3, and Lemma 4.8 will imply the desired $\mathcal{P} = \mathcal{P}'$.

For the claim about X^0 , first of all, the image of i contains infinitely many distinct divisors D_λ for $\lambda \in \Lambda'$, so the Chevalley constructibility theorem (EGA IV₁, 1.8.4) implies that i is dominant. We then let

$$X^0 \subset X \setminus \text{Base}(\mathcal{P})$$

be a nonempty open over which i is flat (EGA IV₃, 11.2.6 (ii)) and let x range over the closed points of $D_\lambda \cap X^0$ for $\lambda \in \Lambda'$. Suppose that for such an $x \in D_\lambda \cap X^0$ the fiber $i^{-1}(x)$ is not a singleton. Then the divisor $i^{-1}(D_\lambda \cap X^0) \subset i^{-1}(X^0)$ meets some π -fibers E_c over closed points $c \in C$ such that E_c is different from the π -fiber D_λ . By construction, the effective divisors E_c on X are all algebraically equivalent to D_λ , so, since the latter

is irreducible, the nonempty intersections $E_c \cap D_\lambda$ have pure codimension 2 in X . This means that the intersections $i^{-1}(D_\lambda \cap X^0) \cap E_c$ are nowhere dense in E_c .

Since $i^{-1}(D_\lambda \cap X^0) \subset i^{-1}(X^0)$ is of pure codimension 1, another application of the Chevalley constructibility theorem then shows that the map $i^{-1}(D_\lambda \cap X^0) \rightarrow C$ given by π is dominant and its image contains a nonempty open $C^0 \subset C$. However, this is impossible: by construction of C , there is a closed point $c \in C^0$ such that $E_c = D_{\lambda'}$ for some $\lambda' \in \Lambda' \setminus \{\lambda\}$, and $i^{-1}(D_\lambda \cap X^0)$ cannot meet E_c because $D_\lambda \cap D_{\lambda'}$ lies in $\text{Base}(\mathcal{P})$, which does not meet X^0 . Thus, $i^{-1}(x)$ is indeed a singleton, as desired. \square

Corollary 4.10. — *If k is uncountable (equivalently, if $|X|$ is uncountable), then every topological pencil on X is algebraic.*

Proof. — To see the parenthetical equivalence it suffices to note that, by Noether normalization, $|X|$ is uncountable if and only if $|\mathbb{A}_k^{\dim(X)}|$ is uncountable. Suppose that $|X|$ is uncountable, let $\mathcal{P} = \{D_\lambda\}_{\lambda \in \Lambda}$ be a topological pencil on X , and fix a $\lambda \in \Lambda$ and a closed point $x \in D_\lambda \setminus \text{Base}(\mathcal{P})$. Since $\text{Base}(\mathcal{P})$ is of codimension ≥ 2 in X , by cutting D_λ by sufficiently general hyperplanes passing through x supplied by the Bertini theorem, we may build an irreducible curve $C \subset X$ that properly meets D_λ but does not meet $\text{Base}(\mathcal{P})$. Then C meets each $D_{\lambda'}$ in finitely many points, to the effect that, since $|C|$ is uncountable and the $D_{\lambda'}$ cover the closed points of X , the set Λ is also uncountable. On the other hand, the Néron–Severi group $\text{NS}(X)$ is countable (SGA 6, XIII, Thm. 5.1). Thus, there is an infinite subset $\Lambda' \subset \Lambda$ such that the $D_{\lambda'}$ for $\lambda' \in \Lambda'$ are pairwise algebraically equivalent, and so also pairwise numerically equivalent (SGA 6, XIII, Thm. 4.6). Proposition 4.9 then shows that the pencil \mathcal{P} is algebraic. \square

5. RECOVERING LINEAR EQUIVALENCE OF DIVISORS

NOTATION. *In this section, we let X be a geometrically normal, geometrically integral, positive-dimensional projective variety over an infinite field k .*

Lemma 5.1. — *For the subgroup $\mathcal{R} \subset \text{Div}(X)$ generated by the differences of linearly equivalent reduced divisors, every class in $\text{Div}(X)/\mathcal{R}$ is represented by a difference of reduced divisors. In particular, \mathcal{R} is the subgroup of all divisors linearly equivalent to 0.*

Proof. — The last assertion follows from the rest because the difference of reduced divisors lies in \mathcal{R} if and only if this difference is linearly equivalent to 0. For the rest, every divisor is a sum of irreducible divisors (with multiplicities), so it suffices to show that every irreducible divisor $D \subset X$ is linearly equivalent to $D_1 - D_2$ for some reduced divisors $D_i \subset X$ that share no irreducible components with divisors $D' \subset X$ in some fixed finite set containing D : then D and $D_1 - D_2$ will agree in $\text{Div}(X)/\mathcal{R}$ and, by iteratively applying this with D' ranging over the D_i from preceding steps, we will represent every class in $\text{Div}(X)/\mathcal{R}$ by a difference of reduced divisors.

To find a desired linear equivalence $D \sim D_1 - D_2$, we first choose a very ample divisor $H \subset X$ and an $m > 0$ such that $D + mH$ is also very ample (HARTSHORNE, 1977, II, Exercise 7.5). Since k is infinite, the Bertini theorem (JOUANOLOU, 1983, 6.10) then supplies our desired geometrically reduced divisors D_i with $D_1 \sim D + mH$ and $D_2 \sim mH$. \square

As for linear equivalence of reduced divisors, the following is the key criterion.

Proposition 5.2 (KOLLÁR, 2020, Theorem 115). — *Suppose that k is perfect and $\dim X \geq 3$, and let $H \subset X$ be an irreducible ample divisor. Reduced divisors $D_1, D_2 \subset X$ are linearly equivalent if and only if for every large enough closed subset $\Sigma \subset X$ of codimension ≥ 2 and every integral curve $C \subset X$ not in $D_1 \cup D_2 \cup \Sigma$, there are*

- irreducible divisors $D', E' \subset X$ not in $D_1 \cup D_2$ with E' ample and containing C ;
- algebraic topological pencils \mathcal{P}_1 and \mathcal{P}_2 with $C \subset \text{Base}(\mathcal{P}_i)$;
- an irreducible ample divisor $E \subset X$ with $(D_i + D' + E') \setminus (E \cup \Sigma)$ connected;

such that

- $D_i + D' + E'$ and E lie in \mathcal{P}_i (so E also contains C);
- all but finitely many closed points of C lie in the k -smooth loci E^{sm} and E'^{sm} ;
- $D_i + D' + E'$ and E lie in the subset of those $D \in \mathcal{P}_i$ for which the function $D \mapsto D \cdot H^{\dim X - 1}$ takes the minimal value attained infinitely many times on \mathcal{P}_i .

Sketch of proof. — We begin with the simpler ‘only if’ direction, suppose that D_1 and D_2 are linearly equivalent, and choose Σ to contain $X \setminus X^{\text{sm}}$. We then choose an irreducible ample divisor D' that shares no components with D_1 and D_2 , does not contain C , and is such that the $\mathcal{O}(D_i) \otimes_{\mathcal{O}_X} \mathcal{O}(D')$ are generated by global sections. This makes the reduced divisors $D_i + D'$ basepoint free, so also Cartier. Moreover, since k is perfect, the integral curve C is generically k -smooth. We may then use the Bertini theorem (KLEIMAN and ALTMAN, 1979, Thm. 7) (which uses the assumption on $\dim(X)$) to find an irreducible ample divisor E' that contains C , is k -smooth at the generic point of C , properly meets every irreducible component of $D_i + D'$, does not contain any irreducible component of Σ , and is such that the $D_i + D' + E'$ are very ample (HARTSHORNE, 1977, II, Exercise 7.5 (d)). Granted that we make sure (as we may) that E' is sufficiently ample, we may then apply the Bertini theorem (KLEIMAN and ALTMAN, 1979, Thm. 7) again, this time with the ample line bundle

$$\mathcal{O}(D_1 + D' + E') \simeq \mathcal{O}(D_2 + D' + E'),$$

to find an irreducible ample divisor $E \subset X$ with $E \sim D_i + D' + E'$ such that E contains C , is k -smooth at the generic point of C , and does not contain any irreducible component of $D_i + D' + E'$ nor any irreducible component of the intersections between E' and the irreducible components of the $D_i + D'$. The

$$(D_i + D' + E') \setminus (E \cup \Sigma)$$

are then connected, and we let \mathcal{P}_i be the linear pencil spanned by $D_i + D' + E'$ and E as in Example 4.5. By construction, these D' , E' , \mathcal{P}_i , and E satisfy all the requirements (to check the requirement about the intersection numbers, one uses the fact that both $D_i + D' + E'$ and E are k -fibers of the pencil and argues with a complete H -intersection curve Y analogous to the one used in the proof of Proposition 4.9).

For the 'if' direction, we may assume that Σ contains $X \setminus X^{\text{sm}}$ and use the Bertini theorem (JOUANOLOU, 1983, Thm. 6.10) to choose C to be a generically smooth, geometrically integral complete H -intersection. With these choices, by Lemma 4.6 1), the pencils \mathcal{P}_i are rational and, by its part 3), then they are even linear. At this point one uses the condition that involves the degree function

$$D \mapsto D \cdot H^{\dim X - 1}$$

to check that both $D_i + D' + E'$ and E are k -fibers of the pencil \mathcal{P}_i (we omit the details of this step here). It then follows from Example 4.5 that $D_i + D' + E' \sim E$, so that also $D_1 \sim D_2$. \square

Corollary 5.3 (KOLLÁR, 2020, Theorem 116). — *Suppose that $\text{char}(k) = 0$ and $\dim X \geq 4$. The topological space $|X|$ determines linear equivalence of divisors on X .*

Sketch of proof. — By Lemma 5.1, it suffices to show that $|X|$ determines linear equivalence between reduced divisors. For this, we explain why the notions and conditions that appear in the linear equivalence criterion of Proposition 5.2 are determined by $|X|$:

- Ampleness of irreducible divisors by Proposition 2.2.
- Algebraicity of topological pencils by Proposition 4.9.
- The function $D \mapsto D \cdot H^{\dim X - 1}$ up to constant multiple by Corollary 3.7.
- All but finitely many closed points of C lying in E^{sm} and E'^{sm} by Proposition 3.5 2): to apply it, we choose Σ to contain $X \setminus X^{\text{sm}}$ (then all but finitely many closed points of C lie in X^{sm}) and we note that E and E' are geometrically connected by the Lefschetz hyperplane theorem (SGA 2_{new}, XII, 3.5). \square

In conclusion, we have now described the second reconstruction step promised in §1:

$$(|X|, \sim_s \text{ of irreducible ample divisors}) \rightsquigarrow (|X|, \sim \text{ of effective divisors}).$$

6. RECOVERING THE PROJECTIVE VARIETY ITSELF

NOTATION. *In this section, we let X be a normal, geometrically integral projective variety of dimension ≥ 2 over an infinite field k .*

The remaining step

$$(|X|, \sim \text{ of effective divisors}) \rightsquigarrow X$$

is a result of LIEBLICH and OLSSON (2019). In this step, the ultimate source of reconstruction is the fundamental theorem of projective geometry that characterizes projective spaces combinatorially in terms of properties of incidence between their points and lines—in effect, for projective spaces this theorem reconstructs the full structure of an algebraic variety from axiomatic combinatorial data. It is fascinating that the collinearity relation for points in projective space encodes rich scheme-theoretic structure. This is not quite unexpected, however—after all, one knows that this relation is capable of encoding, for instance, polynomial equations with integer coefficients (LAFFORGUE, 2003, p. iv, Remarques).

The precise version of such a theorem that Lieblich and Olsson use is as follows, established via an argument that is close to E. Artin's classical proof.

Theorem 6.1 (LIEBLICH and OLSSON, 2019, Theorem 4.4). — *Let V and V' be finite-dimensional vector spaces over infinite fields k and k' , respectively, and let*

$$U \subset \mathrm{Gr}_1(\mathbb{P}(V))(k) \quad \text{and} \quad U' \subset \mathrm{Gr}_1(\mathbb{P}(V'))(k')$$

be collections of lines in $\mathbb{P}(V)$ and $\mathbb{P}(V')$ given by the sets of k -points and k' -points of some nonempty Zariski open subsets of the indicated Grassmannians. For any bijection

$$b: \mathbb{P}(V)(k) \xrightarrow{\sim} \mathbb{P}(V')(k') \quad \text{that induces an inclusion} \quad U \hookrightarrow U',$$

there are a field isomorphism $\iota: k \xrightarrow{\sim} k'$ and a ι -semilinear isomorphism $V \xrightarrow{\sim} V'$ such that the induced isomorphism $\mathbb{P}(V) \xrightarrow{\sim} \mathbb{P}(V')$ agrees with b on some Zariski open containing all the lines in U .

With this theorem in hand, the strategy is to apply it with

$$V := \Gamma(X, \mathcal{O}(mH)) \quad \text{and} \quad V' := \Gamma(X', \mathcal{O}(mH')) \quad \text{for all} \quad m > 0$$

for a suitable very ample divisor $H \subset X$ and a homeomorphic to X projective variety X' that one seeks to show to be isomorphic to X (with H' being the image of H).

Indeed, the isomorphisms $V \xrightarrow{\sim} V'$ then give isomorphisms between the graded components of the coordinate rings that appear in the projective embeddings

$$\begin{aligned} X &\cong \mathrm{Proj} \left(\bigoplus_{m \geq 0} \Gamma(X, \mathcal{O}(mH)) \right) \hookrightarrow \mathrm{Proj} \left(\bigoplus_{m \geq 0} \Gamma(X, \mathcal{O}(H))^{\otimes m} \right), \\ X' &\cong \mathrm{Proj} \left(\bigoplus_{m \geq 0} \Gamma(X', \mathcal{O}(mH')) \right) \hookrightarrow \mathrm{Proj} \left(\bigoplus_{m \geq 0} \Gamma(X, \mathcal{O}(H'))^{\otimes m} \right). \end{aligned}$$

With the help of some additional arguments to make sure that the isomorphisms of the graded pieces are compatible as m varies, one obtains the desired $X \xrightarrow{\sim} X'$.

In view of this strategy, the key becomes defining the subsets of lines U and U' with which to apply Theorem 6.1. Since k is infinite, a line $\ell \subset \mathbb{P}(V)$ is spanned by any two of its k -points, which correspond to k^\times -orbits of sections of $\Gamma(X, \mathcal{O}(mH))$, that is, to effective divisors on X linearly equivalent to mH . The *base locus* of ℓ is the locus of common vanishing on X of these effective divisors. Lieblich and Olsson choose U to consist of all the lines that satisfy the following definition (and analogously for U').

Definition 6.2. With $V := \Gamma(X, \mathcal{O}(mH))$ as above, a line $\ell \subset \mathbb{P}(V)$ is *strongly definable* if there is a closed subset $Z \subset |X|$ of codimension ≥ 2 such that ℓ consists precisely of those effective divisors on X linearly equivalent to mH that contain Z .

The following description of the set of all strongly definable lines is critical for applying Theorem 6.1 to reconstruct the projective variety X .

Lemma 6.3 (LIEBLICH and OLSSON, 2019, Lemma 5.13). — *If the linear system determined by the ample divisor $H \subset X$ is basepoint free, then the strongly definable lines in $\mathbb{P}(V)$ comprise the set of k -points of a nonempty Zariski open of the Grassmannian $\mathrm{Gr}_1(\mathbb{P}(V))$.*

The precise formulation of the reconstruction result that Lieblich and Olsson obtain by carrying out the strategy that we sketched in this section is as follows.

Theorem 6.4 (LIEBLICH and OLSSON, 2019, Proposition 6.2.5). — *For normal, geometrically integral, proper varieties X and X' over infinite fields k and k' , respectively, if X is projective and $\dim X \geq 2$, then any homeomorphism $|X| \xrightarrow{\sim} |X'|$ that matches the linear equivalence relations on effective divisors on X and X' underlies a scheme isomorphism $X \xrightarrow{\sim} X'$.*

This result achieves the final reconstruction step

$$(|X|, \sim \text{ of effective divisors}) \rightsquigarrow X,$$

and hence completes our sketch of the proof of Theorem 1.1.

Acknowledgements. I thank the organizers of the Séminaire Bourbaki for their kind invitation. I thank János Kollár for patiently answering numerous questions. I thank Viện Toán Học for an invitation to give lectures related to the material presented here.

REFERENCES

- GROTHENDIECK, Alexander (2005). *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. French. Vol. 4. Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Séminaire de Géométrie Algébrique du Bois Marie, 1962; Augmenté d'un exposé de Michèle Raynaud. [With an exposé by Michèle Raynaud]; With a preface and edited by Yves Laszlo; Revised reprint of the 1968 French original. Société Mathématique de France, Paris, pp. x+208. ISBN: 2-85629-169-4 (henceforth cited as: SGA 2_{new}).
- BERTHELOT, Pierre, GROTHENDIECK, Alexander, and ILLUSIE, Luc (1971). *Théorie des intersections et théorème de Riemann-Roch. Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967 (SGA 6)*. French. Lecture Notes in Mathematics, Vol. 225. Avec la collaboration de D. Ferrand, J. P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J. P. Serre. Springer-Verlag, Berlin-New York, pp. xii+700 (henceforth cited as: SGA 6).

- BHATT, Bhargav (2012). « Annihilating the cohomology of group schemes », *Algebra Number Theory* **6** (7), pp. 1561–1577.
- BOGOMOLOV, Fedor A., PIRUTKA, Alena, and SILBERSTEIN, Aaron Michael (2016). « Families of disjoint divisors on varieties », *Eur. J. Math.* **2** (4), pp. 917–928.
- GROTHENDIECK, Alexander and DIEUDONNÉ, Jean (1961). « Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I », *Inst. Hautes Études Sci. Publ. Math.* **11**, p. 167 (henceforth cited as: EGA III₁).
- (1964). « Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I », French. *Inst. Hautes Études Sci. Publ. Math.* **20**, p. 259 (henceforth cited as: EGA IV₁).
- (1965). « Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II », French. *Inst. Hautes Études Sci. Publ. Math.* **24**, p. 231 (henceforth cited as: EGA IV₂).
- (1966). « Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III », *Inst. Hautes Études Sci. Publ. Math.* **28**, p. 255 (henceforth cited as: EGA IV₃).
- HARTSHORNE, Robin (1977). *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, pp. xvi+496. ISBN: 0-387-90244-9.
- HOCHSTER, Melvin (1969). « Prime ideal structure in commutative rings », *Trans. Amer. Math. Soc.* **142**, pp. 43–60.
- DE JONG, Johan and al. (2008–). *The Stacks Project*. Available at <http://stacks.math.columbia.edu> (henceforth cited as: SP).
- JOUANOLOU, Jean-Pierre (1983). *Théorèmes de Bertini et applications*. French. Vol. 42. Progress in Math. Birkhäuser Boston, Inc., Boston, MA, pp. ii+127. ISBN: 0-8176-3164-X.
- KLEIMAN, Steven L. (1966). « Toward a numerical theory of ampleness », *Ann. of Math.* (2) **84**, pp. 293–344.
- KLEIMAN, Steven L. and ALTMAN, Allen B. (1979). « Bertini theorems for hypersurface sections containing a subscheme », *Comm. Algebra* **7** (8), pp. 775–790.
- KOLLÁR, János (1996). *Rational curves on algebraic varieties*. Vol. 32. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, pp. viii+320. ISBN: 3-540-60168-6.
- (2020). « What determines a variety? », *preprint*. Available at <https://arxiv.org/abs/2002.12424v2>.
- KOLLÁR, János, LIEBLICH, Max, OLSSON, Martin, and SAWIN, Will (2020). « Topological reconstruction theorems for varieties », *preprint*. Available at <https://arxiv.org/abs/2003.04847v3>.

- LAFFORGUE, Laurent (2003). *Chirurgie des grassmanniennes*. French. Vol. 19. CRM Monograph Series. American Mathematical Society, Providence, RI, pp. xx+170. ISBN: 0-8218-3358-8.
- LIEBLICH, Max and OLSSON, Martin (2019). « A reconstruction theorem for varieties », *preprint*. Available at <https://arxiv.org/abs/1902.04668v1>.
- NÉRON, André (1952). « Problèmes arithmétiques et géométriques rattachés à la notion de rang d'une courbe algébrique dans un corps », French. *Bull. Soc. Math. France* **80**, pp. 101–166.
- POONEN, Bjorn (2001). « Points having the same residue field as their image under a morphism », *J. Algebra* **243** (1), pp. 224–227.
- (2017). *Rational points on varieties*. Vol. 186. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, pp. xv+337. ISBN: 978-1-4704-3773-2.
- RAVINDRA, Girivaru V. and SRINIVAS, Vasudevan (2006). « The Grothendieck–Lefschetz theorem for normal projective varieties », *J. Algebraic Geom.* **15** (3), pp. 563–590.
- WIEGAND, Roger and KRAUTER, William (1981). « Projective surfaces over a finite field », *Proc. Amer. Math. Soc.* **83** (2), pp. 233–237.

Kęstutis Česnavičius

CNRS, Université Paris-Saclay
Laboratoire de mathématiques d'Orsay
F-91405, Orsay, France
E-mail : kestutis@math.u-psud.fr